

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number  
**WO 01/64679 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 487/04**,  
A61K 31/519

**TAGGART, John, J.** [US/US]; 412 Westview Road,  
Elkins Park, PA 19027 (US).

(21) International Application Number: PCT/US01/06688

(74) **Agents: DINNER, Dara, L.** et al.; SmithKline Beecham  
Corporation, Corporate Intellectual Property, UW2220,  
709 Swedeland Road, P.O. Box 1539, King of Prussia, PA  
19406-0939 (US).

(22) International Filing Date: 2 March 2001 (02.03.2001)

(25) Filing Language: English

(81) **Designated States (national):** AE, AL, AU, BA, BB, BG,  
BR, BZ, CA, CN, CO, CZ, DZ, EE, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG,  
MK, MN, MX, MZ, NO, NZ, PL, RO, SG, SI, SK, SL, TR,  
TT, TZ, UA, US, UZ, VN, YU, ZA.

(26) Publication Language: English

(30) **Priority Data:**  
60/186,419 2 March 2000 (02.03.2000) US

(84) **Designated States (regional):** ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) **Applicant (for all designated States except US):**  
**SMITHKLINE BEECHAM CORPORATION**  
[US/US]; One Franklin Plaza, Philadelphia, PA 19103  
(US).

(72) **Inventors; and**

**Published:**

(75) **Inventors/Applicants (for US only): ADAMS, Jerry,**  
**L.** [US/US]; 611 Forest Road, Wayne, PA 19087 (US).  
**BOEHM, Jeffrey, C.** [US/US]; 248 Anthony Road, King  
of Prussia, PA 19406 (US). **HALL, Ralph, F.** [US/US];  
1311 Prospect Hill Road, Villanova, PA 19085 (US).

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) **Title:** 1,5-DISUBSTITUTED-3,4-DIHYDRO-1H-PYRIMIDO[4,5-D]PYRIMIDIN-2-ONE COMPOUNDS AND THEIR USE IN TREATING CSBP/P38 KINASE MEDIATED DISEASES

(57) **Abstract:** Novel substituted pyrimido[4,5-d]pyrimidin-2-one compounds and compositions for use in therapy as CSBP/p38 kinase inhibitors.



**WO 01/64679 A1**

**1,5-DISUBSTITUTED-3,4-DIHYDRO-1H-PYRIMIDO[4,5-D]PYRIMIDIN-2-ONE COMPOUNDS AND THEIR USE IN TREATING CSBP/P38 KINASE MEDIATED DISEASES.****FIELD OF THE INVENTION**

5 This invention relates to a novel group of 1,5-disubstituted-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one compounds, processes for the preparation thereof, the use thereof in treating CSBP/p38 kinase mediated diseases and pharmaceutical compositions for use in such therapy.

**BACKGROUND OF THE INVENTION**

10 Intracellular signal transduction is the means by which cells respond to extracellular stimuli. Regardless of the nature of the cell surface receptor (e. g. protein tyrosine kinase or seven-transmembrane G-protein coupled), protein kinases and phosphatases along with phospholipases are the essential machinery by which the signal is further transmitted within the cell [Marshall, J. C. Cell , **80**, 179-278 (1995)].

15 Protein kinases can be categorized into five classes with the two major classes being, tyrosine kinases and serine / threonine kinases depending upon whether the enzyme phosphorylates its substrate(s) on specific tyrosine(s) or serine / threonine(s) residues [Hunter, T., Methods in Enzymology (Protein Kinase Classification) p. 3, Hunter, T.; Sefton, B. M.; eds. vol. 200, Academic Press; San Diego, 1991].

20 For most biological responses, multiple intracellular kinases are involved and an individual kinase can be involved in more than one signaling event. These kinases are often cytosolic and can translocate to the nucleus or the ribosomes where they can affect transcriptional and translational events, respectively. The involvement of kinases in transcriptional control is presently much better understood than their effect  
25 on translation as illustrated by the studies on growth factor induced signal transduction involving MAP/ERK kinase [Marshall, C. J. Cell , **80**, 179 (1995); Herskowitz, I. Cell , **80**, 187 (1995); Hunter, T. Cell , **80**, 225 (1995); Seger, R., and Krebs, E. G. FASEB J., 726-735 (1995)].

While many signaling pathways are part of cell homeostasis, numerous  
30 cytokines (e.g., IL-1 and TNF) and certain other mediators of inflammation (e.g., COX-2, and iNOS) are produced only as a response to stress signals such as bacterial lipopolysaccharide (LPS). The first indications suggesting that the signal transduction pathway leading to LPS-induced cytokine biosynthesis involved protein kinases came from studies of Weinstein [Weinstein, *et al.*, J. Immunol. **151**, 3829(1993)] but the  
35 specific protein kinases involved were not identified. Working from a similar perspective, Han [Han, *et al.*, Science **265**, 808(1994)] identified murine p38 as a kinase which is tyrosine phosphorylated in response to LPS. Definitive proof of the

involvement of the p38 kinase in LPS-stimulated signal transduction pathway leading to the initiation of proinflammatory cytokine biosynthesis was provided by the independent discovery of p38 kinase by Lee [Lee; *et al.*, Nature, **372**, 739(1994)] as the molecular target for a novel class of anti-inflammatory agents. The discovery of p38 (termed by Lee as CSBP 1 and 2) provided a mechanism of action of a class of anti-inflammatory compounds for which SK&F 86002 was the prototypic example. These compounds inhibited IL-1 and TNF synthesis in human monocytes at concentrations in the low uM range [Lee, *et al.*, Int. J. Immunopharmac. 10(7), 835(1988)] and exhibited activity in animal models which are refractory to cyclooxygenase inhibitors [Lee; *et al.*, Annals N. Y. Acad. Sci., **696**, 149(1993)].

It is now firmly established that CSBP/p38 is a one of several kinases involved in a stress-response signal transduction pathway, which is parallel to and largely independent of the analogous mitogen-activated protein kinase (MAP) kinase cascade. Stress signals, including LPS, pro-inflammatory cytokines, oxidants, UV light and osmotic stress, activate kinases upstream from CSBP/p38 which in turn phosphorylate CSBP/p38 at threonine 180 and tyrosine 182 resulting in CSBP/p38 activation. MAPKAP kinase-2 and MAPKAP kinase-3 have been identified as downstream substrates of CSBP/p38 which in turn phosphorylate heat shock protein Hsp 27 (Figure 1). Additional downstream substrates known to be phosphorylated by p38 include kinases (Mnk1/2, MSK1/2 and PRAK) and transcription factors (CHOP, MEF2, ATF2 and CREB). While many of the signaling pathways required for cytokine biosynthesis remain unknown it appears clear that many of the substrates for p38 listed above are involved. [Cohen, P. Trends Cell Biol., 353-361(1997) and Lee, J. C. *et al.*, Pharmacol. Ther. vol. 82, nos. 2-3, pp. 389-397, 1999].

What is known, however, is that in addition to inhibiting IL-1 and TNF, CSBP/p38 kinase inhibitors (SK&F 86002 and SB 203580) also decrease the synthesis of a wide variety of pro-inflammatory proteins including, IL-6, IL-8, GM-CSF and COX-2. Inhibitors of CSBP/p38 kinase have also been shown to suppress the TNF-induced expression of VCAM-1 on endothelial cells, the TNF-induced phosphorylation and activation of cytosolic PLA2 and the IL-1-stimulated synthesis of collagenase and stromelysin. These and additional data demonstrate that CSBP/p38 is involved not only cytokine synthesis, but also in cytokine signaling [CSBP/P38 kinase reviewed in Cohen, P. Trends Cell Biol., 353-361(1997)].

Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are biological substances produced by a variety of cells, such as monocytes or macrophages. IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions such as

inflammation [See, e.g., Dinarello et al., Rev. Infect. Disease, **6**, 51 (1984)]. The myriad of known biological activities of IL-1 include the activation of T helper cells, induction of fever, stimulation of prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase proteins and the suppression of plasma iron levels.

There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis, and acute synovitis. Recent evidence also links IL-1 activity to diabetes and pancreatic  $\beta$  cells [review of the biological activities which have been attributed to IL-1 Dinarello, *J. Clinical Immunology*, **5** (5), 287-297 (1985)].

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia, secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, or pyresis.

Interleukin-8 (IL-8) is a chemotactic factor produced by several cell types including mononuclear cells, fibroblasts, endothelial cells, and keratinocytes. Its production from endothelial cells is induced by IL-1, TNF, or lipopolysachharide (LPS). IL-8 stimulates a number of functions in vitro. It has been shown to have chemoattractant properties for neutrophils, T-lymphocytes, and basophils. In addition it induces histamine release from basophils from both normal and atopic individuals as well as lysozomal enzyme release and respiratory burst from neutrophils. IL-8 has also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils without de novo protein synthesis, this may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many diseases are characterized by massive neutrophil infiltration. Conditions associated with an increased in IL-8 production (which is responsible for chemotaxis of neutrophil into the inflammatory site) would

benefit by compounds, which are suppressive of IL-8 production.

IL-1 and TNF affect a wide variety of cells and tissues and these cytokines as well as other leukocyte derived cytokines are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these  
5 cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

Inhibition of signal transduction via CSBP/p38, which in addition to IL-1, TNF and IL-8 described above is also required for the synthesis and/or action of several additional pro-inflammatory proteins (i.e., IL-6, GM-CSF, COX-2, collagenase and  
10 stromelysin), is expected to be a highly effective mechanism for regulating the excessive and destructive activation of the immune system. This expectation is supported by the potent and diverse anti-inflammatory activities described for CSBP/p38 kinase inhibitors [Badger, *et al.*, *J. Pharm. Exp. Thera.* **279** (3): 1453-1461.(1996); Griswold, *et al.*, *Pharmacol. Comm.* **7**, 323-229 (1996)].

The field of patents and patent applications disclosing compounds useful for the treatment of p38 mediated diseases has expanded rapidly over the last several years. In most instances the central core molecule has been an imidazole, oxazole or pyrazole derivative, such as those disclosed in WO 93/14081; WO 93/14082; WO 95/02591; WO 95/13067; WO 95/31451; WO 99/58523; WO 98/56377; WO 97/16442; WO  
20 99/57101; WO 00/39116; and WO 00/31063. Newer ring systems include cycloalkenyl, pyrimidine, pyrazine, and triazole cores such as WO 00/25791; WO 98/24782; WO 99/17776; WO 00/10563; WO 00/25791; and WO 00/35911; and multi ring systems, such as WO 99/64400; WO 98/22457; WO 00/20402; WO 00/12497; WO 99/61426 and WO 99/58502.

However, despite all of this research effort there still remains a need for  
25 treatments in this field, for compounds which inhibit the CSBP/p38/RK kinase, and are useful in the treatment of disease mediated thereby.

#### BRIEF DESCRIPTION OF THE DRAWINGS

30 Figure 1 demonstrates the p38 kinase cascade.

Figure 2 demonstrates a Citric Acid Induced Cough Model.

Figure 3 demonstrates an Antigen- or LTD4 – Induced Hypertussive Model in the  
Guinea Pig


Figure 4 demonstrates Effects of Dextromethorphan or Codeine On Citric Acid-  
35 Induced Cough in Guinea Pigs.

#### SUMMARY OF THE INVENTION

This invention relates to the novel compounds of Formulas (I) to (V) and (Ia) to (Va), and pharmaceutical compositions comprising a compound of Formula (I) to (V) and (Ia) to (Va), and a pharmaceutically acceptable diluent or carrier.

This invention relates to a method of treating, including prophylaxis, of a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formulas (I) to (V) and (Ia) to (Va).

Accordingly, the present invention provides a compound of Formula (I):



10

wherein

R<sub>1</sub> is an aryl or heteroaryl ring, which ring is optionally substituted;

R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub> alkyl, aryl, arylC<sub>1-10</sub>alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub>alkyl moiety; and wherein each of these moieties,

15

excluding hydrogen, are optionally substituted;

R<sub>3</sub> is a C<sub>1-10</sub> alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub> alkyl, arylC<sub>1-10</sub> alkyl, heteroarylC<sub>1-10</sub> alkyl, or heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are optionally substituted;

20

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl, or an optionally substituted arylC<sub>1-4</sub>alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members, which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may be optionally substituted;

25

R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, may be optionally substituted;

30

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted


aryl or optionally substituted arylC<sub>1-4</sub> alkyl;

Z is oxygen or sulfur;

or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention is the novel compounds of Formula

5 (II) represented by the structure:



wherein

R<sub>1</sub> is an aryl or a heteroaryl ring, which ring may be optionally substituted;

10 R<sub>2</sub> is hydrogen, C<sub>1-10</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, are optionally substituted;

15 R<sub>3</sub> is an C<sub>1-10</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, aryl, arylC<sub>1-10</sub>alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are optionally substituted;

Y is CR<sub>b</sub>, C(O), N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>;

R<sub>b</sub> is hydrogen, C<sub>1-2</sub> alkyl, NR<sub>c</sub>, hydroxy, thio, C<sub>1-2</sub> alkoxy, or S(O)<sub>m</sub>C<sub>1-2</sub> alkyl;

20 R<sub>c</sub> is hydrogen or C<sub>1-2</sub> alkyl;

R<sub>d</sub> is hydrogen or C<sub>1-2</sub> alkyl;

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub>, (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

m is 0 or an integer having a value of 1 or 2;

n is 0 or an integer having a value of 1 to 10;

25 R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl, or an optionally substituted arylC<sub>1-4</sub>alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may be optionally substituted;


30 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl; and wherein each of

these moieties, excluding hydrogen, may be optionally substituted;  
 R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub> or optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;


Z is oxygen or sulfur;

5 or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention are the novel compounds of Formula (III) represented by the structure:



(III) or



(IIIa)

wherein

R<sub>1</sub> is an aryl, or a heteroaryl ring, which ring may be optionally substituted;

R<sub>2</sub> is a heterocyclic or heterocyclic C<sub>1-10</sub> alkyl ring, which is ring is optionally substituted;

15 R<sub>2</sub>' is a heterocyclic, heterocyclic C<sub>1-10</sub> alkyl, aryl, or heteroaryl ring, which ring is optionally substituted;

R<sub>3</sub> is an aryl or heteroaryl ring, which ring is optionally substituted;

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>; provided that when X is (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, then R<sub>4</sub>R<sub>14</sub> must form the optionally

20 substituted cyclized ring optionally comprising an oxygen, sulfur or nitrogen moiety;

n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

25 R<sub>4</sub> and R<sub>14</sub> are independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl, or optionally substituted aryl-C<sub>1-4</sub> alkyl; or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring is optionally substituted;

30 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, aryl-C<sub>1-10</sub> alkyl, heteroaryl or a heteroaryl-C<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, is optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted




aryl, or optionally substituted arylC<sub>1-4</sub> alkyl;

Z is oxygen or sulfur;

or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention are the novel compounds of Formula  
5 (IV) represented by the structures:



wherein

R<sub>1</sub> is an aryl or a heteroaryl ring, which ring is optionally substituted;

R<sub>2</sub> is hydrogen, C<sub>1-10</sub>alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub>alkyl, aryl,  
10 arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a  
heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties,  
excluding hydrogen, are optionally substituted;

R<sub>3</sub> is a C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub>alkyl, aryl,  
arylC<sub>1-10</sub>alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a  
15 heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are  
optionally substituted;

Y is a bond, CR<sub>b</sub>, C(O), N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>;

R<sub>b</sub> is hydrogen, C<sub>1-2</sub>alkyl, NR<sub>c</sub>, hydroxy, thio, C<sub>1-2</sub>alkoxy, S(O)<sub>m</sub>C<sub>1-2</sub>alkyl;

R<sub>c</sub> is hydrogen or C<sub>1-2</sub>alkyl;

20 R<sub>d</sub> is hydrogen or C<sub>1-2</sub>alkyl;

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted  
25 C<sub>1-4</sub> alkyl, optionally substituted aryl, or optionally substituted arylC<sub>1-4</sub> alkyl,  
or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a  
heterocyclic ring of 5 to 7 members which ring optionally contains an additional  
heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring is optionally  
substituted;


30 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl,  
aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety, and wherein  
each of these moieties, excluding hydrogen, is optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or an optionally substituted aryl-C<sub>1-4</sub> alkyl;


Z is oxygen or sulfur;

or a pharmaceutically acceptable salt thereof.

5 Accordingly, the present invention also provides for a compound of Formula (V):



(V) or



(Va)

wherein

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

10 n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

R<sub>1</sub> is an optionally substituted aryl ring;

R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub> alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a

15 heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen are optionally substituted;

R<sub>3</sub> is an optionally substituted aryl ring;

R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl, or optionally substituted aryl-C<sub>1-4</sub> alkyl,

20 or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring is optionally substituted;

25 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, may be optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or an optionally substituted arylC<sub>1-4</sub> alkyl;

Z is oxygen or sulfur;

30 or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to the novel compounds of Formulas (I) to (V), and Formula (Ia) to (Va), or a pharmaceutically acceptable salt thereof; pharmaceutical compositions comprising a compound of Formula (I) to (V), and (Ia) to (Va), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent thereof; and also to the use of these compounds in the treatment or prophylaxis of CSBP mediated diseases in a mammal in need thereof.

As will be readily recognized, the difference between compounds of Formula (I) to (V) and (Ia) to (Va) lie in the unsaturation of the pyrimidine-2-one ring. The respective R<sub>1</sub>, R<sub>2</sub>, X and R<sub>3</sub> terms are the same for both groups within a formula set, i.e. I and Ia. For purposes herein, everything applicable to Formula (I) is also applicable to Formula (Ia), Formula (II) and (IIa), Formula (III) and (IIIa), and Formula (V) and (Va) unless otherwise indicated. It should be noted that compounds of Formula (I), (II), (III) and (V) are all subsets of compounds of Formula (IV).

More specifically, the present invention is directed to the novel compounds of Formulas (I) and (Ia), or pharmaceutically acceptable salts thereof.

Suitably, for these compounds of Formula (I) and (Ia), R<sub>1</sub> is an aryl, or a heteroaryl ring, which ring is optionally substituted. The ring may be substituted one or more times, in any ring, suitably 1 to 4 times, preferably 1 to 2 times, independently, by halogen, C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, cyano, nitro, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)OR<sub>8</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>a</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(O)H, S(O)<sub>m</sub>R<sub>5</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>8</sub>, ZC(Z)R<sub>11</sub>, NR<sub>10</sub>C(Z)R<sub>11</sub>, or NR<sub>10</sub>S(O)<sub>2</sub>R<sub>7</sub>. Preferably the ring is an aryl ring, more preferably phenyl, which ring is unsubstituted or substituted one or more times by halogen, preferably chloro or fluoro, or alkyl, preferably methyl. Suitably, the ring is substituted in the 4-position, and if di-substituted, it is in the 2-,4- position of the phenyl ring.

Suitably m is 0 or an integer having a value of 1 or 2.

Suitably, v is 0 or an integer having a value of 1 or 2.

Suitably, Z is oxygen or sulfur.

Suitably, R<sub>a</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl,

heterocyclalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted aryl, or an optionally substituted aryl-C<sub>1-6</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen to which they are attached in NR<sub>4</sub>R<sub>14</sub> substituent may form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may be optionally substituted.

Suitably, R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>4</sub>R<sub>14</sub>, excluding the moieties SR<sub>5</sub> (wherein m is 0) being SNR<sub>4</sub>R<sub>14</sub>, S(O)<sub>2</sub>R<sub>5</sub> being SO<sub>2</sub>H (wherein m is 2) and S(O)R<sub>5</sub> being SOH, (wherein m is 1).

Suitably, R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclalkyl, heterocyclalkyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety, and wherein each these moieties may be optionally substituted. It is recognized herein that hydrogen can not be substituted and is therefore excluded.

Suitably, R<sub>7</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclalkylC<sub>1-6</sub> alkyl, heteroaryl, or a heteroarylC<sub>1-6</sub>alkyl moiety; and wherein each of these moieties may be optionally substituted.

Suitably, R<sub>8</sub> is hydrogen, C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclalkyl, heterocyclalkylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclalkyl, heterocyclalkyl alkyl, heteroaryl, and heteroaryl alkyl moieties, may be optionally substituted.

Suitably, t is an integer having a value of 1 to 3.

Suitably, R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl, or optionally substituted aryl-C<sub>1-4</sub> alkyl.

Suitably, R<sub>10</sub> and R<sub>20</sub> are each independently selected from hydrogen or C<sub>1-4</sub> alkyl.

Suitably, R<sub>11</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub>alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclalkyl, heterocyclalkylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclalkyl, heterocyclalkylalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, R<sub>3</sub> is a C<sub>1-10</sub>alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub>alkyl, arylC<sub>1-10</sub> alkyl, heteroarylC<sub>1-10</sub> alkyl, or a heterocyclylC<sub>1-10</sub> alkyl moiety, and wherein each of these moieties may be optionally substituted, independently, one or more times, suitably 1 to 4 times, preferably 1 to 2 times, (in any applicable ring) with C<sub>1-10</sub> alkyl, halosubstituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub>cycloalkenylC<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, R<sub>3</sub> is an optionally substituted C<sub>3-7</sub> cycloalkyl, or arylC<sub>1-10</sub> alkyl, more preferably optionally substituted cyclohexyl, benzyl or phenethyl. Preferably, the ring is unsubstituted or substituted one or more times, independently with halogen, more preferably fluorine, or chlorine, or alkyl, such as methyl. The ring is preferably substituted in the 2-position or di-substituted in the 2,6- position, more preferably with fluorine, methyl, difluoro, or dimethyl.

Suitably, n is 0, or an integer having a value of 1 to 10.

Suitably, R<sub>2</sub> is hydrogen, C<sub>1-10</sub>alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein these moieties, excluding hydrogen, may be optionally substituted one or more times, suitably 1 to 4 times, independently (in any applicable ring) with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub>cycloalkenylC<sub>1-10</sub>alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxyalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety.

Suitably, X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>.

Suitably, when X is the group OR<sub>2</sub>, then R<sub>2</sub> is preferably an optionally substituted C<sub>1-10</sub> alkyl, an optionally substituted aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic or heterocyclyl moiety. Preferably, when the R<sub>2</sub>

moiety is an alkyl it is a substituted or unsubstituted C<sub>1-6</sub> alkyl. Suitably, the alkyl substituents are as defined herein, but are preferably, halogen, hydroxy, or NR<sub>4</sub>R<sub>14</sub>. When R<sub>2</sub> is an optionally substituted heterocyclic, or heterocyclic alkyl ring, it is preferably a pyrrolidine, piperidine, piperazine, morpholine ring or a pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or a morpholinyl alkyl. More preferably, it is a piperidine or an alkyl substituted piperidine, such as N-methyl piperidine.

Preferably, when X is the group S(O)<sub>m</sub>R<sub>2</sub>, R<sub>2</sub> is a substituted or unsubstituted alkyl, aryl or aryl alkyl.

Preferably, when X is the group (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, n is preferably 0 or 1, more preferably 0. Preferably, if the R<sub>4</sub> and R<sub>14</sub> moiety cyclize the ring is a 5 or 6 membered ring, such as a pyrrolidine, piperidine, piperazine, or morpholine containing ring, which ring may be optionally substituted. Preferably when R<sub>4</sub> or R<sub>14</sub> is an optionally substituted C<sub>1-6</sub>alkyl, which chain may be a straight or branched chain, it is substituted one or more times, suitably 1 to 4 times, preferably 1 or 2 times, with halogen; hydroxy; hydroxy substituted C<sub>1-10</sub>alkyl; C<sub>1-10</sub> alkoxy; halosubstituted C<sub>1-10</sub> alkoxy; S(O)<sub>m</sub> C<sub>1-4</sub>alkyl, wherein m is 0, 1 or 2; NR<sub>4</sub>R<sub>14</sub>, such as amino or mono or -disubstituted C<sub>1-6</sub> alkyl or wherein the R<sub>4</sub>R<sub>14</sub> can cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S; C<sub>3-7</sub>cycloalkyl; C<sub>3-7</sub>cycloalkyl C<sub>1-10</sub> alkyl group; halosubstituted C<sub>1-10</sub> alkyl, such CF<sub>2</sub>CF<sub>2</sub>H, or CF<sub>3</sub>; optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl; and wherein these aryl or arylalkyl moieties may also themselves be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl; C<sub>1-10</sub> alkoxy; S(O)<sub>m</sub> C<sub>1-4</sub> alkyl; amino, mono & di-substituted C<sub>1-6</sub> alkyl amino, such as in the NR<sub>4</sub>R<sub>14</sub> group; C<sub>1-4</sub> alkyl, or CF<sub>3</sub>.

Preferably, the alkyl chain is branched, such as in t-butyl or isopropyl. More preferably the alkyl chain is substituted one or more times, independently by halogen, hydroxy, amino, or mono or di C<sub>1-4</sub> alkyl substituted amino.

Preferably, when X is the group (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>, n is preferably 0 or 1, more preferably 0. The R<sub>2</sub> moiety is preferably, hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heterocyclic, substituted or unsubstituted heterocyclic alkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heteroarylalkyl. Such groups preferably include imidazole, tetrazole, pyrrolidine, piperidine, piperazine, morpholine, pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or morpholinyl alkyl.

Preferably, when X is the group R<sub>2</sub>, then R<sub>2</sub> is preferably an optionally substituted heteroaryl, such as an imidazole, or tetrazole, or is an optionally substituted heterocyclic ring, such as a pyrrolidine, piperidine, piperazine, or a morpholine ring

5 Another aspect of the present invention is directed to the compounds of Formula (II) and (IIa).

Suitably, for compounds of Formula (II) and (IIa), R<sub>1</sub> is an aryl or a heteroaryl ring, which ring may be optionally substituted. The ring may be substituted one or more times, in any ring, suitably 1 to 4 times, preferably 1 to 2  
10 times, independently, by halogen, C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, cyano, nitro, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)OR<sub>8</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>a</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(O)H, S(O)<sub>m</sub>R<sub>5</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>8</sub>, ZC(Z)R<sub>11</sub>, NR<sub>10</sub>C(Z)R<sub>11</sub>, or NR<sub>10</sub>S(O)<sub>2</sub>R<sub>7</sub>. Preferably the ring is an aryl ring, more preferably phenyl, which ring is unsubstituted or substituted one or more times by  
15 halogen, preferably chloro or fluoro, or alkyl, preferably methyl. Suitably, the ring is substituted in the 4-position, and if di-substituted, it is in the 2-,4- position of the phenyl ring.

Suitably, v is 0 or an integer having a value of 1 or 2.

Suitably, Z is oxygen or sulfur.

20 Suitably, R<sub>a</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylalkyl, heterocyclyl,  
25 heterocyclylalkyl, heteroaryl, or heteroarylalkyl moiety may be optionally substituted.

Suitably, R<sub>4</sub> and R<sub>14</sub> is each independently selected from hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted aryl, or optionally substituted aryl-C<sub>1-6</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are  
30 attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may be optionally substituted.

Suitably, R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>4</sub>R<sub>14</sub>, excluding the moieties SR<sub>5</sub> (wherein m is 0) being SNR<sub>4</sub>R<sub>14</sub>, S(O)<sub>2</sub>R<sub>5</sub>  
35 being SO<sub>2</sub>H (wherein m is 2) and S(O)R<sub>5</sub> being SOH, (wherein m is 1).

Suitably, R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl

moiety, and wherein each of these moieties may be optionally substituted (excluding hydrogen).

Suitably, R<sub>7</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub>alkyl, heteroaryl, or a heteroarylC<sub>1-6</sub>alkyl; and wherein each of these moieties may  
5 be optionally substituted.

Suitably, R<sub>8</sub> is hydrogen, C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or  
10 (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, t is an integer having a value of 1 to 3.

Suitably, R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl,  
15 optionally substituted aryl, or optionally substituted aryl-C<sub>1-4</sub> alkyl.

Suitably, R<sub>10</sub> and R<sub>20</sub> are each independently selected from hydrogen or C<sub>1-4</sub> alkyl.

Suitably, R<sub>11</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub>alkynyl, C<sub>3-7</sub>cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>,  
20 (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

25 Suitably m is 0 or an integer having a value of 1 or 2.

Suitably, X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>.

Suitably, Y is CR<sub>b</sub>, C(O), N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>.

Suitably, R<sub>b</sub> is hydrogen, C<sub>1-2</sub> alkyl, NR<sub>c</sub>, hydroxy, thio, C<sub>1-2</sub> alkoxy, S(O)<sub>m</sub>C<sub>1-2</sub> alkyl.  
30

Suitably, R<sub>c</sub> is hydrogen or C<sub>1-2</sub> alkyl.

Suitably, R<sub>d</sub> is hydrogen or C<sub>1-2</sub> alkyl.

Suitably, R<sub>3</sub> is a C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety, and wherein each of these moieties may be  
35 optionally substituted, independently, one or more times, suitably 1 to 4 times, preferably 1 to 2 times, (in any applicable ring) with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>



alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>  
 S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>,  
 5 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, R<sub>3</sub> is an optionally substituted C<sub>3-7</sub> cycloalkyl, aryl, or arylC<sub>1-10</sub>  
 alkyl, more preferably optionally substituted cyclohexyl, phenyl, benzyl or  
 10 phenethyl. Preferably, the ring is unsubstituted or substituted one or more times,  
 independently with halogen, more preferably fluorine, or chlorine, or alkyl, such as  
 methyl. The ring is preferably substituted in the 2-position or di-substituted in the 2-  
 ,6- position, more preferably with fluorine, methyl, difluoro, or dimethyl.

Suitably, n is 0, or an integer having a value of 1 to 10.

15 Suitably, R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl,  
 aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a  
 heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen,  
 may be optionally substituted one or more times, suitably 1 to 4 times, independently (in  
 any applicable ring), with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub>  
 20 alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub>  
 cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>,  
 25 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxyalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-,  
 wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety.

Suitably, when X is the group OR<sub>2</sub>, then R<sub>2</sub> is preferably an optionally  
 30 substituted C<sub>1-10</sub> alkyl, an optionally substituted aryl, arylalkyl, heteroaryl,  
 heteroarylalkyl, heterocyclic or heterocyclicalkyl moiety. Preferably, when the R<sub>2</sub>  
 moiety is an alkyl it is a substituted or unsubstituted C<sub>1-6</sub> alkyl moiety as defined  
 above. Preferably, the alkyl substituents are independently selected from halogen,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>.

35 When R<sub>2</sub> is an optionally substituted heterocyclic or heterocyclic alkyl ring,  
 it is preferably a pyrrolidine, piperidine, piperazine, or morpholine ring, or a  
 pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or a morpholinyl alkyl ring.

More preferably, it is a piperidine or an alkyl substituted piperidine, such as N-methyl piperidine.

Preferred substituents include halogen,  $C_{1-10}$ alkyl, halosubstituted  $C_{1-10}$ alkyl,  $(CR_{10}R_{20})_nNR_4R_{14}$ ,  $(CR_{10}R_{20})_nOR_6$ , and  $(CR_{10}R_{20})_nC(Z)OR_6$ .

5 Preferably, when X is the group  $S(O)_mR_2$ ,  $R_2$  is a substituted or unsubstituted  $C_{1-10}$ alkyl, aryl or arylalkyl.

Preferably, when X is the group  $(CH_2)_nNR_4R_{14}$ , n is preferably 0 or 1, more preferably 0. Preferably if the  $R_4$  and  $R_{14}$  moiety cyclize the ring is a 5 or 6  
10 membered ring, such as a pyrrolidine, piperidine, piperazine, or morpholine containing ring, which ring may be optionally substituted. Preferably, when  $R_4$  or  $R_{14}$  is an optionally substituted  $C_{1-6}$ alkyl, which chain may be a straight or branched chain, it is substituted one or more times, suitably 1 to 4 times, preferably 1  
or 2 times, with halogen; hydroxy; hydroxy substituted  $C_{1-10}$ alkyl;  $C_{1-10}$ alkoxy; halosubstituted  $C_{1-10}$ alkoxy;  $S(O)_mC_{1-4}$ alkyl, wherein m is 0, 1 or 2;  $NR_4R_{14}$ ,  
15 such as amino or mono or -disubstituted  $C_{1-6}$ alkyl or wherein the  $R_4R_{14}$  can cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S;  $C_{3-7}$ cycloalkyl;  $C_{3-7}$ cycloalkyl  $C_{1-10}$ alkyl group; halosubstituted  $C_{1-10}$ alkyl, such as  $CF_2CF_2H$ , or  $CF_3$ ; optionally substituted aryl, such as phenyl, or an  
20 optionally substituted arylalkyl, such as benzyl or phenethyl; and wherein these aryl or arylalkyl moieties may also themselves be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl;  $C_{1-10}$ alkoxy;  $S(O)_mC_{1-4}$ alkyl; amino, mono & di-substituted  $C_{1-6}$ alkyl amino, such as in the  $NR_4R_{14}$  group;  $C_{1-4}$ alkyl, or  $CF_3$ .

25 Preferably, the alkyl chain is branched, such as in t-butyl or isopropyl. More preferably the alkyl chain is substituted one or more times, independently by halogen, hydroxy, amino, or mono or di  $C_{1-4}$ alkyl substituted amino.

Preferably, when X is the group  $(CH_2)_nNR_2R_4$ , n is preferably 0 or 1, more preferably 0. The  $R_2$  moiety is preferably, hydrogen, substituted or unsubstituted  
30 alkyl, substituted or unsubstituted heterocyclic, substituted or unsubstituted heterocyclic alkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heteroarylalkyl. Such groups preferably include imidazole, tetrazole, pyrrolidine, piperidine, piperazine, morpholine, and 2-,3- or 4-pyridyl, pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or morpholinyl alkyl.

35 Preferably, when X is the group  $R_2$ , then  $R_2$  is preferably an optionally substituted heteroaryl, such as an imidazole, or tetrazole, or is an optionally substituted heterocyclic ring, such as a pyrrolidine, piperidine, piperazine, or a

morpholine ring

Another aspect of the present invention are the novel compounds of Formula (III) and (IIIa).

Suitably, for compounds of Formula (III) and (IIIa), R<sub>1</sub> is an aryl or  
 5 heteroaryl ring, which ring may be optionally substituted. The ring may be substituted one or more times, in any ring, suitably 1 to 4 times, preferably 1 to 4 times, independently, by halogen, C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, cyano, nitro, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)OR<sub>8</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>a</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(O)H, S(O)<sub>m</sub>R<sub>5</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>8</sub>, ZC(Z)R<sub>11</sub>,  
 10 NR<sub>10</sub>C(Z)R<sub>11</sub>, or NR<sub>10</sub>S(O)<sub>2</sub>R<sub>7</sub>. Preferably the ring is an aryl ring, more preferably phenyl, which ring is unsubstituted or substituted one or more times by halogen, preferably chloro or fluoro, or alkyl, preferably methyl. Suitably, the ring is substituted in the 4-position, and if di-substituted, it is in the 2-,4- position of the phenyl ring.

15 Suitably m is 0 or an integer having a value of 1 or 2.

Suitably, v is 0 or an integer having a value of 1 or 2.

Suitably, Z is oxygen or sulfur.

Suitably, R<sub>a</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl,  
 20 heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl moieties may be optionally substituted.

25 Suitably, R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted aryl, or an optionally substituted aryl-C<sub>1-6</sub> alkyl; or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may  
 30 be optionally substituted.

Suitably, R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>4</sub>R<sub>14</sub>, excluding the moieties SR<sub>5</sub> (wherein m is 0) being SNR<sub>4</sub>R<sub>14</sub>, S(O)<sub>2</sub>R<sub>5</sub> being SO<sub>2</sub>H (wherein m is 2) and S(O)R<sub>5</sub> being SOH, (wherein m is 1).

Suitably, R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl,  
 35 heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, may be optionally substituted.

Suitably, R<sub>7</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub>alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl moiety; and wherein each of these moieties may be optionally substituted.

Suitably, R<sub>8</sub> is hydrogen, C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub>alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroaryl alkyl moieties, may be optionally substituted.

Suitably, t is an integer having a value of 1 to 3.

Suitably, R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl.

Suitably, R<sub>10</sub> and R<sub>20</sub> are each independently selected from hydrogen or C<sub>1-4</sub> alkyl.

Suitably, R<sub>11</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>; provided that when X is (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, then R<sub>4</sub>R<sub>14</sub> forms the optionally substituted cyclized ring optionally comprising an oxygen, sulfur or nitrogen moiety as defined herein.

Suitably, R<sub>3</sub> is an optionally substituted aryl, or optionally substituted heteroaryl moiety. The R<sub>3</sub> moieties be optionally substituted one or more times independently with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub> S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, R<sub>3</sub> is an optionally substituted aryl, more preferably optionally substituted phenyl. Preferably, the ring is unsubstituted or substituted one or more times, independently with halogen, more preferably fluorine, or chlorine, or alkyl, such as methyl. The ring is preferably substituted in the 2-position or di-substituted in the 2-,6- position, more preferably with fluorine, methyl, difluoro, or dimethyl.

Suitably, n is 0, or an integer having a value of 1 to 10. Suitably, R<sub>2</sub> is a heterocyclic, or heterocyclylC<sub>1-10</sub> alkyl ring, which ring or rings(s) are optionally substituted one or more times, suitably 1 to 4, independently with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxyalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety. Suitably, R<sub>2</sub>' is a heterocyclic, heterocyclylC<sub>1-10</sub> alkyl, aryl, or heteroaryl ring, which ring or rings(s) are optionally substituted one or more times, suitably 1 to 4 times, in any ring, independently with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxyalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety.

Suitably, the optionally substituted R<sub>2</sub> or R<sub>2</sub>' heterocyclic ring is a pyrrolidine, piperidine, piperazine, or morpholine ring or a pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or a morpholinyl alkyl. More preferably, it is a piperidine or an alkyl substituted piperidine, such as N-methyl piperidine.

Suitably, when R<sub>2</sub>' is an optionally substituted aryl or heteroaryl ring, the ring is an optionally substituted phenyl, imidazole or tetrazole ring.

Another aspect of the present invention is the novel compounds of Formula (IV) and (IVa).

Suitably, for compounds of Formula (IV) and (IVa), R<sub>1</sub> is an aryl or heteroaryl ring, which ring is optionally substituted. The ring may be substituted one or more times, in any ring, suitably 1 to 4 times, preferably 1 to 2 times, independently, by halogen, C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, cyano, nitro,  
5 (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(Z)OR<sub>8</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>a</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>C(O)H, S(O)<sub>m</sub>R<sub>5</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>8</sub>, ZC(Z)R<sub>11</sub>, NR<sub>10</sub>C(Z)R<sub>11</sub>, or NR<sub>10</sub>S(O)<sub>2</sub>R<sub>7</sub>. Preferably the ring is an aryl ring, more preferably phenyl, which ring is unsubstituted or substituted one or more times by halogen, preferably chloro or fluoro, or alkyl, preferably methyl. Suitably, the ring is  
10 substituted in the 4-position, and if di-substituted, it is in the 2-,4- position of the phenyl ring. Most preferably R<sub>1</sub> is a 2-methyl-4-fluoro phenyl.

Suitably m is 0 or an integer having a value of 1 or 2.

Suitably, v is 0 or an integer having a value of 1 or 2.

Suitably, Z is oxygen or sulfur.

15 Suitably, R<sub>a</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl,  
20 heterocyclylalkyl, heteroaryl, and heteroarylalkyl moieties may be optionally substituted.

Suitably, R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted aryl or an optionally substituted aryl-C<sub>1-6</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are  
25 attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring may be optionally substituted.

Suitably, R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>4</sub>R<sub>14</sub>, excluding the moieties SR<sub>5</sub> (wherein m is 0) being SNR<sub>4</sub>R<sub>14</sub>, S(O)<sub>2</sub>R<sub>5</sub>  
30 being SO<sub>2</sub>H (wherein m is 2) and S(O)R<sub>5</sub> being SOH, (wherein m is 1).

Suitably, R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl; and wherein each of these moieties, excluding hydrogen, may be optionally  
substituted.

35 Suitably, R<sub>7</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl; and wherein each of these moieties may be optionally substituted.

Suitably, R<sub>8</sub> is hydrogen, C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or  
 5 (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclyl alkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, t is an integer having a value of 1 to 3.

Suitably, R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl,  
 10 optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl.

Suitably, R<sub>10</sub> and R<sub>20</sub> are each independently selected from hydrogen or C<sub>1-4</sub> alkyl.

Suitably, R<sub>11</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl,  
 15 heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, R<sub>3</sub> is hydrogen, an C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these  
 20 moieties, excluding hydrogen, may be optionally substituted, independently, one or more times, suitably 1 to 4 times, preferably 1 to 2 times, in any ring if applicable, with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>,  
 25 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, R<sub>3</sub> is an optionally substituted C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub>alkyl, or a cycloalkyl moiety, more preferably optionally substituted C<sub>1-5</sub> alkyl,  
 35 cyclohexyl, phenyl, benzyl or phenethyl. Preferably, the ring is unsubstituted or substituted one or more times, independently with halogen, more preferably fluorine, or chlorine, or alkyl, such as methyl. The ring is preferably substituted in the 2-

position or di-substituted in the 2-,6- position, more preferably with fluorine, methyl, difluoro, or dimethyl.

Suitably, X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub>, (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>.

Suitably, R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclicC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, may optionally substituted one or more times, suitable 1 to 4 times, independently, and in any ring, with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub> S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety.

Suitably, n is 0, or an integer having a value of 1 to 10.

Suitably, when X is the group OR<sub>2</sub>, then R<sub>2</sub> is preferably an optionally substituted C<sub>1-10</sub> alkyl, an optionally substituted aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic or heterocyclicalkyl moiety. Preferably, when the R<sub>2</sub> moiety is an alkyl it is a substituted or unsubstituted C<sub>1-6</sub> alkyl. Suitably, the alkyl substituents are halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub> S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, the alkyl substituents are halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>.

When R<sub>2</sub> is an optionally substituted heterocyclic or heterocyclic alkyl ring, it is preferably a pyrrolidine, piperidine, piperazine, or morpholine ring, or is a pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or morpholinyl alkyl ring. More preferably, it is a piperidine or an alkyl substituted piperidine, such as N-methyl piperidine.

Preferred substituents include halogen, C<sub>1-10</sub>alkyl, halosubstituted C<sub>1-10</sub>alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, and (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>.



Preferably when X is the group  $S(O)_mR_2$ ,  $R_2$  is a substituted or unsubstituted  $C_{1-10}$ alkyl, aryl or arylalkyl. While the X group  $S(O)_mR_2$  moiety is a compound of Formula (IV) it is also a key intermediate in the process of making other compounds of Formula (IV) as can be seen by Scheme I herein. Preferably for this use, m is 0 and  $R_2$  is a short chain alkyl, such as methyl.

Preferably, when X is the group  $(CH_2)_nNR_4R_{14}$ , n is preferably 0 or 1, more preferably 0. Preferably, if the  $R_4$  and  $R_{14}$  moiety cyclize the ring is a 5 or 6 membered ring, such as a pyrrolidine, piperidine, piperazine, or morpholine containing ring, which ring may be optionally substituted. Preferably, when  $R_4$  or  $R_{14}$  is an optionally substituted  $C_{1-6}$ alkyl, which chain may be a straight or branched chain, it is substituted one or more times, independently, suitably 1 to 4 times, preferably 1 or 2 times, with halogen; hydroxy; hydroxy substituted  $C_{1-10}$ alkyl;  $C_{1-10}$  alkoxy; halosubstituted  $C_{1-10}$  alkoxy;  $S(O)_mC_{1-4}$ alkyl, wherein m is 0, 1 or 2;  $NR_4R_{14}$ , such as amino or mono or -disubstituted  $C_{1-4}$  alkyl or wherein the  $R_4R_{14}$  can cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S;  $C_{3-7}$ cycloalkyl;  $C_{3-7}$ cycloalkyl  $C_{1-10}$  alkyl group; halosubstituted  $C_{1-10}$  alkyl, such  $CF_2CF_2H$ , or  $CF_3$ ; optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl; and wherein these aryl or arylalkyl moieties may also themselves be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl;  $C_{1-10}$  alkoxy;  $S(O)_mC_{1-4}$  alkyl; amino, mono & di-substituted  $C_{1-4}$  alkyl amino, such as in the  $NR_4R_{14}$  group;  $C_{1-4}$  alkyl, or  $CF_3$ .

Preferably, the alkyl chain is branched, such as in t-butyl or isopropyl. More preferably the alkyl chain is substituted one or more times, independently by halogen, hydroxy, amino, or mono or di  $C_{1-4}$  alkyl substituted amino.

Preferably, when X is the group  $(CH_2)_nNR_2R_4$ , n is preferably 0 or 1, more preferably 0. Preferably one of  $R_2$  or  $R_4$  is hydrogen and the other is an optionally substituted moiety. The  $R_2$  moiety is preferably, hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heterocyclic, substituted or unsubstituted heterocyclic alkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heteroarylalkyl. When  $R_2$  is an optionally substituted heterocyclic, heterocyclic alkyl, or heteroaryl group such groups are preferably an imidazole, tetrazole, pyrrolidine, piperidine, piperazine, morpholine and 2-, 3- or 4-pyridyl; pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or morpholinyl alkyl.

When the  $R_2$  moiety is an optionally substituted  $C_{1-10}$  alkyl, the chain may be straight or branched, and substituted one or more times, suitably 1 to 4 times,

independently by halogen,  $(\text{CR}_{10}\text{R}_{20})_n\text{OR}_6$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{SH}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{S(O)}_m\text{R}_7$ ,  
 $(\text{CR}_{10}\text{R}_{20})_n\text{NHS(O)}_2\text{R}_7$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{CN}$ ,  
 $(\text{CR}_{10}\text{R}_{20})_n\text{S(O)}_2\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_nC(\text{Z})\text{R}_6$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{OC(Z)}\text{R}_6$ ,  
 $(\text{CR}_{10}\text{R}_{20})_nC(\text{Z})\text{OR}_6$ ,  $(\text{CR}_{10}\text{R}_{20})_nC(\text{Z})\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_{10}\text{C(Z)}\text{R}_6$ ,  
5  $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_{10}\text{C(=NR}_{10})\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{OC(Z)}\text{NR}_4\text{R}_{14}$ ,  
 $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_{10}\text{C(Z)}\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_{10}\text{C(Z)}\text{OR}_7$ , or dioxyalkylene group of  
the formula  $-\text{O}-(\text{CH}_2)_s-\text{O}-$ , wherein  $s$  is 1 to 3, preferably  $s$  is 2 yielding a 1,3-  
dioxyethylene moiety.

Preferably, when  $X$  is the group  $\text{R}_2$ , then  $\text{R}_2$  is preferably an optionally  
10 substituted heteroaryl, such as an imidazole, or tetrazole, or is an optionally  
substituted heterocyclic ring, such as a pyrrolidine, piperidine, piperazine, or  
morpholine ring.

Suitably,  $Y$  is a bond,  $\text{CR}_b$ ,  $\text{C(O)}$ ,  $\text{N(R}_d)$ , oxygen, or  $\text{S(O)}_m$ . Preferably  $Y$  is  
a bond.

15 Suitably,  $\text{R}_b$  is hydrogen,  $\text{C}_{1-2}$  alkyl,  $\text{NR}_c$ , hydroxy, thio,  $\text{C}_{1-2}$  alkoxy,  
 $\text{S(O)}_m\text{C}_{1-2}$  alkyl.

Suitably,  $\text{R}_c$  is hydrogen or  $\text{C}_{1-2}$  alkyl.

Suitably,  $\text{R}_d$  is hydrogen or  $\text{C}_{1-2}$  alkyl.

Suitably,  $m$  is 0 or an integer having a value of 1 or 2.

20 Another aspect of the present invention is the novel compounds of Formula  
(V) and (Va).

Suitably, for compounds of Formula (V) and (Va),  $\text{R}_1$  is an optionally  
substituted aryl ring, preferably a phenyl ring. The phenyl or naphthyl ring or ring(s)  
may be substituted one or more times, in any ring, preferably 1 to 4 times,  
25 independently, by substituents selected from halogen,  $\text{C}_{1-4}$  alkyl, halo-substituted-  
 $\text{C}_{1-4}$  alkyl, cyano, nitro,  $(\text{CR}_{10}\text{R}_{20})_v\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_v\text{C(Z)}\text{NR}_4\text{R}_{14}$ ,  
 $(\text{CR}_{10}\text{R}_{20})_v\text{C(Z)}\text{OR}_8$ ,  $(\text{CR}_{10}\text{R}_{20})_v\text{COR}_a$ ,  $(\text{CR}_{10}\text{R}_{20})_v\text{C(O)H}$ ,  $\text{S(O)}_m\text{R}_5$ ,  
 $(\text{CR}_{10}\text{R}_{20})_v\text{OR}_8$ ,  $\text{ZC(Z)}\text{R}_{11}$ ,  $\text{NR}_{10}\text{C(Z)}\text{R}_{11}$ , or  $\text{NR}_{10}\text{S(O)}_2\text{R}_7$ . Preferably the ring  
is unsubstituted or substituted one or more times by halogen, preferably chloro or  
30 fluoro, or alkyl, preferably methyl. Suitably, the ring is substituted in the 4-position,  
and if di-substituted, in the 2-,4- position of the phenyl ring.

Suitably,  $v$  is 0 or an integer having a value of 1 or 2.

Suitably,  $Z$  is oxygen or sulfur.

Suitably,  $\text{R}_a$  is  $\text{C}_{1-4}$  alkyl, halo-substituted  $\text{C}_{1-4}$  alkyl,  $\text{C}_{2-4}$  alkenyl,  $\text{C}_{2-4}$   
35 alkynyl,  $\text{C}_{3-7}$  cycloalkyl,  $\text{C}_{5-7}$  cycloalkenyl, aryl, aryl $\text{C}_{1-4}$  alkyl, heteroaryl,  
heteroaryl $\text{C}_{1-4}$  alkyl, heterocyclyl, heterocyclyl $\text{C}_{1-4}$  alkyl,  $(\text{CR}_{10}\text{R}_{20})_v\text{OR}_7$ ,  
 $(\text{CR}_{10}\text{R}_{20})_v\text{S(O)}_m\text{R}_7$ ,  $(\text{CR}_{10}\text{R}_{20})_v\text{NHS(O)}_2\text{R}_7$ , or  $(\text{CR}_{10}\text{R}_{20})_v\text{NR}_4\text{R}_{14}$ ; and

wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclyl alkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, R<sub>4</sub> and R<sub>14</sub> is each independently selected from hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-6</sub> alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, which ring may be optionally substituted.

Suitably, R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>4</sub>R<sub>14</sub>, excluding the moieties SR<sub>5</sub> (wherein m is 0) being SNR<sub>4</sub>R<sub>14</sub>, S(O)<sub>2</sub>R<sub>5</sub> being SO<sub>2</sub>H (wherein m is 2) and S(O)R<sub>5</sub> being SOH, (wherein m is 1).

Suitably, R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl; and wherein each of these moieties, excluding hydrogen, may be optionally substituted.

Suitably, R<sub>7</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl; and wherein each of these moieties may be optionally substituted.

Suitably, R<sub>8</sub> is hydrogen, C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclyl alkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally substituted.

Suitably, t is an integer having a value of 1 to 3.

Suitably, R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl.

Suitably, R<sub>10</sub> and R<sub>20</sub> are each independently selected from hydrogen or C<sub>1-4</sub> alkyl.

Suitably, R<sub>11</sub> is C<sub>1-4</sub> alkyl, halo-substituted C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclyl, heterocyclylC<sub>1-4</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>OR<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>4</sub>R<sub>14</sub>; and wherein the alkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroaryl alkyl moieties may be optionally

substituted.

Suitably m is 0 or an integer having a value of 1 or 2.

Suitably, R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, may optionally substituted one or more times independently, preferably 1 to 2 times, with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub> S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>, or a dioxalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety.

Suitably, R<sub>3</sub> is an optionally substituted aryl ring, which ring may be optionally substituted one or more times, independently, in any ring, with C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub> alkyl, halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>SH, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>7</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub> S(O)<sub>2</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>6</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>10</sub>) NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OC(Z)NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z) NR<sub>4</sub>R<sub>14</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>7</sub>.

Preferably, R<sub>3</sub> is an optionally substituted phenyl. Preferably, the ring is unsubstituted or substituted one or more times, independently with halogen, more preferably fluorine, or chlorine, or alkyl, such as methyl. The ring is preferably substituted in the 2-position or di-substituted in the 2-,6- position, more preferably with fluorine, methyl, difluoro, or dimethyl.

Suitably, n is 0, or an integer having a value of 1 to 10.

Suitably, X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub>, (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>.

Suitably, when X is the group OR<sub>2</sub>, then R<sub>2</sub> is preferably an optionally substituted C<sub>1-10</sub> alkyl, an optionally substituted aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic or heterocyclicalkyl moiety. Preferably, when the R<sub>2</sub> moiety is an alkyl it is a substituted or unsubstituted C<sub>1-6</sub> alkyl moiety. Preferably, the alkyl substituents are independently selected from halogen, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, or

$(\text{CR}_{10}\text{R}_{20})_n\text{NR}_4\text{R}_{14}$ .

When  $\text{R}_2$  is an optionally substituted heterocyclic or heterocyclic alkyl ring, it is preferably a pyrrolidine, piperidine, piperazine, or morpholine ring, or a pyrrolidinyl alkyl, piperidinyl alkyl, piperazinyl alkyl, or a morpholinyl alkyl ring.

5 More preferably, it is a piperidine or an alkyl substituted piperidine, such as N-methyl piperidine.

Preferred substituents include halogen,  $\text{C}_{1-10}$ alkyl, halosubstituted  $\text{C}_{1-10}$ alkyl,  $(\text{CR}_{10}\text{R}_{20})_n\text{NR}_4\text{R}_{14}$ ,  $(\text{CR}_{10}\text{R}_{20})_n\text{OR}_6$ , and  $(\text{CR}_{10}\text{R}_{20})_n\text{C}(\text{Z})\text{OR}_6$ .

10 Preferably when X is the group  $\text{S}(\text{O})_m\text{R}_2$ , and  $\text{R}_2$  is a substituted or unsubstituted  $\text{C}_{1-10}$ alkyl, aryl or arylalkyl.

Preferably, when X is the group  $(\text{CH}_2)_n\text{NR}_4\text{R}_{14}$ , n is preferably 0 or 1, more preferably 0. Preferably if the  $\text{R}_4$  and  $\text{R}_{14}$  moiety cyclize the ring is a 5 or 6 membered ring, such as a pyrrolidine, piperidine, piperazine, or morpholine containing ring, which ring may be optionally substituted. Preferably, when  $\text{R}_4$  or  
 15  $\text{R}_{14}$  is an optionally substituted  $\text{C}_{1-6}$ alkyl, which chain may be a straight or branched chain, it is substituted one or more times, suitably 1 to 4 times, preferably 1 or 2 times, with halogen; hydroxy; hydroxy substituted  $\text{C}_{1-10}$ alkyl;  $\text{C}_{1-10}$  alkoxy; halosubstituted  $\text{C}_{1-10}$  alkoxy;  $\text{S}(\text{O})_m\text{C}_{1-4}$ alkyl, wherein m is 0, 1 or 2;  $\text{NR}_4\text{R}_{14}$ , such as amino or mono or -disubstituted  $\text{C}_{1-6}$  alkyl or wherein the  $\text{R}_4\text{R}_{14}$  can  
 20 cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S;  $\text{C}_{3-7}$ cycloalkyl;  $\text{C}_{3-7}$ cycloalkyl  $\text{C}_{1-10}$  alkyl group; halosubstituted  $\text{C}_{1-10}$  alkyl, such  $\text{CF}_2\text{CF}_2\text{H}$ , or  $\text{CF}_3$ ; optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl; and wherein these aryl  
 25 or arylalkyl moieties may also themselves be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl;  $\text{C}_{1-10}$  alkoxy;  $\text{S}(\text{O})_m\text{C}_{1-4}$  alkyl; amino, mono & di-substituted  $\text{C}_{1-6}$  alkyl amino, such as in the  $\text{NR}_4\text{R}_{14}$  group;  $\text{C}_{1-4}$  alkyl, or  $\text{CF}_3$ .

30 Preferably, the alkyl chain is branched, such as in t-butyl or isopropyl. More preferably the alkyl chain is substituted one or more times, independently by halogen, hydroxy, amino, or mono or di  $\text{C}_{1-4}$  alkyl substituted amino.

Preferably, when X is the group  $(\text{CH}_2)_n\text{NR}_2\text{R}_4$ , n is preferably 0 or 1, more preferably 0. The  $\text{R}_2$  moiety is preferably, hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heterocyclic, substituted or unsubstituted  
 35 heterocyclic alkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heteroarylalkyl. Such groups preferably include morpholinoethyl, pyrroleethyl, piperidineethyl, pyridylethyl, or piperidine.

Preferably, when X is the group R<sub>2</sub>, then R<sub>2</sub> is preferably an optionally substituted heteroaryl, such as an imidazole, or tetrazole, or an optionally substituted heterocyclic ring, such as a pyrrolidine, piperidine, piperazine, or a morpholine ring

5 As used herein, "optionally substituted" unless specifically defined shall mean, substituent groups which may be substituted on the applicable moiety one or more times, independently, by halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; hydroxy substituted C<sub>1-10</sub>alkyl; C<sub>1-10</sub> alkoxy, such as methoxy or ethoxy; halosubstituted C<sub>1-10</sub> alkoxy; S(O)<sub>m</sub> alkyl, wherein m is 0, 1 or 2, such  
10 as methyl thio, methylsulfinyl or methyl sulfonyl; NR<sub>4</sub>R<sub>14</sub>, such as amino or mono or -disubstituted C<sub>1-4</sub> alkyl or wherein the R<sub>4</sub>R<sub>14</sub> can cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S; C<sub>1-10</sub> alkyl, C<sub>3-7</sub>cycloalkyl, or C<sub>3-7</sub>cycloalkyl C<sub>1-10</sub> alkyl group, such as methyl, ethyl,  
15 propyl, isopropyl, t-butyl, etc. or cyclopropyl methyl; halosubstituted C<sub>1-10</sub> alkyl, such CF<sub>2</sub>CF<sub>2</sub>H, or CF<sub>3</sub>; optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl; and wherein the aryl or arylalkyl moieties may also themselves be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl; C<sub>1-10</sub> alkoxy; S(O)<sub>m</sub>alkyl; amino,  
20 mono & di-substituted C<sub>1-4</sub> alkyl amino, such as in the NR<sub>4</sub>R<sub>14</sub> group; C<sub>1-4</sub> alkyl, or CF<sub>3</sub>.

Suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of inorganic and organic acids, such as hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid,  
25 ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid.

In addition, pharmaceutically acceptable salts of compounds of Formulas (I) to (IV) may also be formed with a pharmaceutically acceptable cation, for instance,  
30 if a substituent group comprises a carboxy moiety. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations.

The term "halo" or "halogens" is used herein to mean the halogens, chloro, fluoro, bromo and iodo.

35 The term "C<sub>1-10</sub>alkyl" or "alkyl" or "alkyl<sub>1-10</sub>" is used herein to mean both straight and branched chain radicals of 1 to 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-

propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.

The term "cycloalkyl" is used herein to mean cyclic radicals, preferably of 3 to 8 carbons, including but not limited to, cyclopropyl, cyclopentyl, cyclohexyl, and the like.

5       The term "cycloalkenyl" is used herein to mean cyclic radicals, preferably of 5 to 8 carbons, which have at least one bond including but not limited to cyclopentenyl, cyclohexenyl, and the like.

      The term "alkenyl" is used herein at all occurrences to mean straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.

      The term "aryl" is used herein to mean phenyl and naphthyl.

      The term "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") is used herein to mean a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, quinoline, isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, thiazole, thiadiazole, tetrazole, triazole, imidazole, or benzimidazole.

20       The term "heterocyclic" (on its own or in any combination, such as "heterocyclylalkyl") is used herein to mean a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydropyran, or imidazolidine.

25       The term "aralkyl" or "heteroarylalkyl" or "heterocyclicalkyl" or "cycloalkyl alkyl" or cycloalkenylalkyl" is used herein to mean a C<sub>1-4</sub> alkyl chain, as defined above, which chain is attached to the respective aryl, heteroaryl, heterocyclic, cycloalkyl or cycloalkenyl moiety as also defined herein unless otherwise indicated.

30       The term "sulfinyl" is used herein to mean the oxide S(O) of the corresponding sulfide; the term "thio" refers to the sulfide; and the term "sulfonyl" refers to the fully oxidized S(O)<sub>2</sub> moiety.

      The term "aroyl" is used herein to mean C(O)Ar, wherein Ar is as phenyl, naphthyl, or an arylalkyl derivative such as defined above, such group include but are not limited to benzyl and phenethyl.

35       The term "alkanoyl" is used herein to mean C(O)C<sub>1-10</sub> alkyl, wherein the alkyl chain is as defined above.

      It is recognized that the compounds of the present invention may exist as

stereoisomers, regioisomers, or diastereoisomers. These compounds may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds are included within the scope of the present invention.

5 Exemplified compounds of the compounds of Formulas (I) to (IV) and (Ia) to (IVa) include those described in the synthetic examples, and those noted below, and their pharmaceutically acceptable salt thereof.

Exemplified compounds of Formula (I), include:

10 7-Methylsulfanyl-5-phenyl-1-(1-phenylethyl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (Ia), include:

7-Methylsulfanyl-5-phenyl-1-(1-phenylethyl)-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (II), include:

15 5-Benzyl-7-(3-morpholin-4-yl-propylamino)-1-phenyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (IIa), include:

5-Benzyl-7-(3-morpholin-4-yl-propylamino)-1-phenyl-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (III), include:

20 7-(4-Methyl-piperazin-1-yl)-1,5-diphenyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (IIIa), include:

7-(4-Methyl-piperazin-1-yl)-1,5-diphenyl-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

Exemplified compounds of Formula (IV), include, but are not limited to,

25 1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfanyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

Exemplified compounds of Formula (IVa), include:

7-Methylsulfanyl-1,5-diphenyl-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one.

30 The compounds of Formula (I), (II), (III), (IV) and (V) may be obtained by applying synthetic procedures, described herein. The synthesis provided for is applicable to producing compounds of Formula (I) to (V) having a variety of different R<sub>1</sub>, R<sub>2</sub>, Y, X, and R<sub>3</sub> groups which are reacted, employing optional substituents which are suitably protected, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, in those cases, then affords compounds of  
35 the nature generally disclosed.

Once the nucleus has been established, further compounds of Formula (I) may be prepared by applying standard techniques for functional group



interconversion, well known in the art. For instance: C(O)NR<sub>4</sub>R<sub>14</sub> from CO<sub>2</sub>CH<sub>3</sub> by heating with or without catalytic metal cyanide, e.g. NaCN, and HNR<sub>4</sub>R<sub>14</sub> in CH<sub>3</sub>OH; OC(O)R<sub>3</sub> from OH with e.g., ClC(O)R<sub>3</sub> in pyridine; NR<sub>10</sub>-C(S)NR<sub>4</sub>R<sub>14</sub> from NHR<sub>10</sub> with an alkylisothiocyanate or thiocyanic acid; NR<sub>10</sub>C(O)OR<sub>7</sub> from  
 5 NHR<sub>10</sub> with the alkyl chloroformate; NR<sub>10</sub>C(O)NR<sub>4</sub>R<sub>14</sub> from NHR<sub>10</sub> by treatment with an isocyanate, e.g. HN=C=O or R<sub>10</sub>N=C=O; NR<sub>10</sub>-C(O)R<sub>7</sub> from NHR<sub>10</sub> by treatment with Cl-C(O)R<sub>7</sub> in pyridine; C(=NR<sub>10</sub>)NR<sub>4</sub>R<sub>14</sub> from C(NR<sub>4</sub>R<sub>14</sub>)SR<sub>3</sub> with H<sub>3</sub>NR<sub>3</sub><sup>+</sup>OAc<sup>-</sup> by heating in alcohol; C(NR<sub>4</sub>R<sub>14</sub>)SR<sub>3</sub> from C(S)NR<sub>4</sub>R<sub>14</sub> with R<sub>6</sub>-I in an inert solvent, e.g. acetone; C(S)NR<sub>4</sub>R<sub>14</sub> (where R<sub>4</sub> or R<sub>14</sub> is not  
 10 hydrogen) from C(S)NH<sub>2</sub> with HNR<sub>4</sub>R<sub>14</sub>-C(=NCN)-NR<sub>4</sub>R<sub>14</sub> from C(=NR<sub>4</sub>R<sub>14</sub>)-SR<sub>3</sub> with NH<sub>2</sub>CN by heating in anhydrous alcohol, alternatively from C(=NH)-NR<sub>4</sub>R<sub>14</sub> by treatment with BrCN and NaOEt in EtOH; NR<sub>10</sub>SO<sub>2</sub>R<sub>3</sub> from NHR<sub>10</sub> by treatment with ClSO<sub>2</sub>R<sub>3</sub> by heating in pyridine; NR<sub>10</sub>C(S)R<sub>6</sub> from NR<sub>10</sub>C(O)R<sub>6</sub> by treatment with Lawesson's reagent [2,4-bis(4-methoxyphenyl)-  
 15 1,3,2,4-dithiadiphosphetane-2,4-disulfide]; NR<sub>10</sub>SO<sub>2</sub>CF<sub>3</sub> from NHR<sub>6</sub> with triflic anhydride and base wherein R<sub>3</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>10</sub>, R<sub>4</sub> and R<sub>14</sub> are as defined in Formula (IV) herein.

Further, other precursors of the various substituent groups herein may be other groups of compounds of Formula (I) to (V) which may be interconverted by  
 20 applying standard techniques for this functional group interconversion. Such as use of the S-alkyl intermediate for X in compounds of Formula (IV) which are oxidized and displaced by a suitable nucleophile to yield other final products of Formula (IV). Also for example, wherein a moiety is a halo substituted C<sub>1-10</sub> alkyl it can be converted to the corresponding C<sub>1-10</sub> alkylN<sub>3</sub> derivative by reacting with a suitable  
 25 azide salt, and thereafter if desired can be reduced to the corresponding C<sub>1-10</sub>alkylNH<sub>2</sub> compound, which in turn can be reacted with R<sub>7</sub>S(O)<sub>2</sub>X wherein X is halo (e.g., chloro) to yield the corresponding C<sub>1-10</sub>alkylNHS(O)<sub>2</sub>R<sub>7</sub> compound.

Alternatively wherein the moiety is a halo-substituted C<sub>1-10</sub>-alkyl it can be reacted with an amine R<sub>4</sub>R<sub>14</sub>NH to yield the corresponding C<sub>1-10</sub>-alkylNR<sub>4</sub>R<sub>14</sub>  
 30 compound, or can be reacted with an alkali metal salt of R<sub>7</sub>SH to yield the corresponding C<sub>1-10</sub>alkylSR<sub>7</sub> compound.


Suitable protecting groups for use with hydroxyl groups and nitrogen groups are well known in the art and described in many references, for instance, Protecting Groups in Organic Synthesis, Greene T W, Wiley-Interscience, New York, 1981.  
 35 Suitable examples of hydroxyl protecting groups include silyl ethers, such as t-butyl dimethyl or t-butyl diphenyl, and alkyl ethers, such as methyl connected by an

alkyl chain of variable link, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>.

Pharmaceutically acid addition salts of compounds of Formula (I) to (V), and (Ia) to (Va) may be obtained in known manner, for example by treatment thereof with an appropriate amount of acid in the presence of a suitable solvent.

5

**Scheme 1**



In Schemes I and II herein, X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl, or heteroaryl group, and Y-R<sub>1</sub> is as defined for Formula (IV) compounds.

10

Commercially available 4,6-dihydroxy-2-methylmercaptopyrimidine (1) (which can also be prepared by the literature methods and utilized as described below, with S-alkyl other than S-methyl or S-aryl) was converted to the nitrile (3) by the literature procedure [see Santilli, et al., *J. Heterocycl. Chem.* **1971**, 8, 445-453] (**Scheme 1**). Compound 3 reacts with one equivalent of amine by procedures

analogous to those described for related compounds, to afford the 2-methylsulfanyl-4-chloro-6-amino-pyrimidine-5-carbonitrile (**4**) [See Tumkevicius, S. *Liebigs Ann.* **1995**, 1703-1705].

The Suzuki reaction of **4** with aryl boronic acids using a palladium catalyst, such as, *tetrakis*(triphenylphosphine) palladium(0) catalyst proceeds in good to excellent yield. Alternatively, the bi-aryl coupling reaction of (**4**) can be performed using aryl or heteroaryl organozinc, organocopper, organotin, or other organometallic reagents known to afford bi-aryl cross-coupling products [See for example Solberg, J.; Undheim, K. *Acta Chemica Scandinavia* **1989**, 62-68]. Displacement of the chlorine in **4** may also be achieved with nitrogen nucleophiles [For related aminations see US patent 3,631,045 and 3,910,913]. Displacement of the chlorine in **4** is also possible with S nucleophiles, [See Tumkevicius, S. *Liebigs Ann.* **1995**, 1703-1705] or O, or alkyl nucleophiles.

The 7-methylsulfamyl-1,5-disubstituted-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-ones (**7**) are prepared by reduction of the nitriles (**5**) preferably with a hydride reducing agent, such as lithium aluminum hydride or NaBH<sub>4</sub>, preferably in an ethereal solvent, such as THF, Et<sub>2</sub>O, glyme or dioxane to produce the di-amine (**6**). In cases where functionality is incompatible with the more reactive hydride reducing agents, alternative, selective, reduction methods for the nitriles include samarium di-iodide in H<sub>3</sub>PO<sub>4</sub>, Raney Ni catalyzed hydrogenation, or sodium borohydride-cobaltous chloride. The resulting diamines **6** can be cyclized to (**7**) with carbonyldiimidazole or a mixed anhydride such as ethyl chloroformate, or for less reactive diamines, with phosgene or triphosgene and a suitable tertiary amine base at temperatures between -78° and 100°C.

Oxidation of the sulfides (**7**) with an oxidant such as one or two equivalents of *meta*-chloroperoxybenzoic acid or Oxone<sup>®</sup> affords either the sulfoxides or sulfones. Oxidation of the sulfides to sulfones can also be effected by OsO<sub>4</sub> and catalytic tertiary amine N-oxide. Other methods for sulfide oxidation include the use of hydrogen peroxide, other peracids, oxygen, ozone, organic peroxides, potassium and zinc permanganate, potassium persulfate, and sodium hypochlorite.

Both 2-pyrimidinyl sulfones and sulfoxides related to compounds of Formula (IV) wherein X is SO-alkyl or SO<sub>2</sub>-alkyl have been reported in the literature to be displaced by a wide variety of nucleophiles. Thus the analogs of Formula (IV) compounds wherein X is an alkyl sulfone or sulfoxide may be displaced by primary and secondary alkylamines without additional base catalysis, preferably in a polar aprotic solvent, such as but not limited to, N-methyl pyrrolidin-2-one (NMP), and at varying temperatures depending upon the nucleophilicity of the amine. For instance

displacement of the sulfone of analogs of Formula (IV) compounds with ethanolamine, in NMP, occurs in 30 min. at about 65° while a more hindered amine such as Tris(hydroxymethyl)aminomethane requires elevated temperatures and extended reaction times (such as, about 80° and about a 24 hour reaction time). The sulfone may also be displaced with substituted arylamine, or heteroaryl amines at elevated temperatures, sometimes requiring formation of the aryl or heteroarylamine anion as with sodium hydride, or other suitable base, in DMSO. In addition, the sulfoxide analogs of Formula (IV) compounds may be readily displaced with aluminum salts of aryl or heteroaryl amines as previously described in the patent literature (WO 9932121).

Likewise, sulfone and sulfoxide analogs of (IV) may be displaced with aryl or heteroaryl or alkyl thiols or alkyl or aryl or heteroaryl alcohols. For instance analogs of IV containing sulfones as the X substituents may be displaced with sodium alkoxide in the alcohol or alternatively reactive alkoxide or phenoxide nucleophiles may be generated from the alcohol or phenol with a suitable base such as NaH or sodium bis-trimethylsilyl amide in a polar aprotic solvent such as DMSO.


Similarly 2-pyrimidinyl sulfones related to (IV) may be displaced with carbon nucleophiles such as aryl or alkyl Grignard reagents or related organometalics such as organo lithium, zinc, tin or boron. These reactions may, in some cases, require transition metal catalysis such as with Pd or Ni catalysts. Displacement of related 2-pyrimidine sulfones with cyanide, malonate anions, unactivated enolates, or heterocyclic C nucleophiles such as 1-methylimidazole anion, by the generation of the anion with NaH or other suitable base in THF also has precedent (see for example, Chem Pharm Bull. **1987**, 4972-4976.). For example analogs of (IV) compounds wherein X is an alkyl sulfone may be displaced with the anion of 1-methyl imidazole, generated by treatment of 1-methyl imidazole with n-butyl lithium in a solvent such as THF at temperatures of about -70°, to afford the C-alkylated product substituted on the imidazole C-2.

The 3,4-Dihydro-1H-pyrimido[4,5-d]pyrimidin-2-ones, unsubstituted at the 7 position (**9**) (**Scheme 1**) can be obtained by Raney Ni hydrogenolysis of the SMe compound (**7**) as well as by direct synthesis from the analog of **1** (in **Scheme 1**) which lacks the S alkyl substituent.

The oxidized series of analogs of Formula (IVa) compounds may be prepared from **5** by partial reduction of the nitrile with a suitable hydride reducing agent such as diisobutylaluminum hydride, preferably at low temperature, and avoiding hydrolytic conditions, to afford the imine (**Scheme 2**). Cyclization of the imine with phosgene or a less reactive equivalent of phosgene such as


carbonyldiimidazole affords **1a** (Scheme 2).

**Scheme 2**




- 5 Substitution at the 7 position of the 3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-ones with aryl substituents is also readily achieved by preforming the 2-aryl pyrimidines as depicted in **Scheme 3** below. While the scheme displays R<sub>1</sub> and X as an aryl moiety, it is merely for representation purposes herein.

**Scheme 3.**




10

Another aspect of the present invention are the intermediate compounds of Formula (A):



wherein Y-R<sub>1</sub> and R<sub>3</sub> are as defined herein for Formula (IV); and X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl or heteroaryl moiety. Preferably X<sub>1</sub> is a C<sub>1-10</sub> alkyl, more preferably a methyl or propyl.

5           Another aspect of the present invention are the intermediate compounds of Formula (C):



wherein Y-R<sub>1</sub> and R<sub>3</sub> are as defined herein for Formula (IV); and X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl or heteroaryl moiety. Preferably X<sub>1</sub> is a C<sub>1-10</sub> alkyl, more preferably a methyl or propyl.

10

## METHODS OF TREATMENT

The compounds of Formula (I) to (V) and (Ia) to (Va) or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated cytokine production by such mammal's cell, such as but not limited to monocytes and/or macrophages; or by the exacerbation or excessive or unregulated production of the CSBP protein.

15

For purposes herein, compounds of Formula (I) to (V) and (Ia) to (Va) will all be referred to as compounds of Formula (I) unless otherwise indicated.

20

Compounds of Formula (I) are capable of inhibiting proinflammatory cytokines, such as IL-1, IL-6, IL-8, and TNF and are therefore of use in therapy. IL-1, IL-6, IL-8 and TNF affect a wide variety of cells and tissues and these cytokines, as well as other leukocyte-derived cytokines, are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these pro-inflammatory cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

25

Accordingly, the present invention provides a method of treating a cytokine-

mediated disease which comprises administering an effective cytokine-interfering amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Compounds of Formula (I) are capable of inhibiting inducible proinflammatory proteins, such as COX-2, also referred to by many other names  
5 such as prostaglandin endoperoxide synthase-2 (PGHS-2) and are therefore of use in therapy. These proinflammatory lipid mediators of the cyclooxygenase (CO) pathway are produced by the inducible COX-2 enzyme. Regulation, therefore of COX-2 which is responsible for the these products derived from arachidonic acid, such as prostaglandins affect a wide variety of cells and tissues are important and  
10 critical inflammatory mediators of a wide variety of disease states and conditions. Expression of COX-1 is not effected by compounds of Formula (I). This selective inhibition of COX-2 may alleviate or spare ulcerogenic liability associated with inhibition of COX-1 thereby inhibiting prostoglandins essential for cytoprotective effects. Thus inhibition of these pro-inflammatory mediators is of benefit in  
15 controlling, reducing and alleviating many of these disease states. Most notably these inflammatory mediators, in particular prostaglandins, have been implicated in pain, such as in the sensitization of pain receptors, or edema. This aspect of pain management therefore includes treatment of neuromuscular pain, headache, cancer pain, and arthritis pain. Compounds of Formula (I) or a pharmaceutically acceptable  
20 salt thereof, are of use in the prophylaxis or therapy in a human, or other mammal, by inhibition of the synthesis of the COX-2 enzyme.

Accordingly, the present invention provides a method of inhibiting the synthesis of COX-2 which comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. The present  
25 invention also provides for a method of prophylaxis treatment in a human, or other mammal, by inhibition of the synthesis of the COX-2 enzyme.

In particular, compounds of Formula (I) or a pharmaceutically acceptable salt thereof are of use in the prophylaxis or therapy of any disease state in a human, or other mammal, which is exacerbated by or caused by excessive or unregulated IL-1,  
30 IL-6, IL-8 or TNF production by such mammal's cell, such as, but not limited to, monocytes and/or macrophages.

Accordingly, in another aspect, this invention relates to a method of inhibiting the production of IL-1 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or  
35 a pharmaceutically acceptable salt thereof.

There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include

rheumatoid arthritis, osteoarthritis, meningitis, ischemic and hemorrhagic stroke, neurotrauma/closed head injury, stroke, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease, tuberculosis, atherosclerosis, muscle degeneration, multiple sclerosis, cachexia, bone resorption, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis. Recent evidence also links IL-1 activity to diabetes, pancreatic  $\beta$  cell diseases and Alzheimer's disease.

Use of a CSAID for the treatment of CSBP mediated disease states, can include, but not be limited to neurodegenerative diseases, such as Alzheimer's disease (as noted above), Parkinson's disease and multiple sclerosis, etc.

In a further aspect, this invention relates to a method of inhibiting the production of TNF in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, chronic pulmonary inflammatory disease and chronic obstructive pulmonary disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, such as osteoporosis, cardiac, brain and renal reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, brain infections including encephalitis (including HIV-induced forms), cerebral malaria, meningitis, ischemic and hemorrhagic stroke, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, inflammatory bowel disease, Crohn's disease, ulcerative colitis and pyresis.

Compounds of Formula (I) are also useful in the treatment of viral infections, where such viruses are sensitive to upregulation by TNF or will elicit TNF production *in vivo*. The viruses contemplated for treatment herein are those that produce TNF as a result of infection, or those which are sensitive to inhibition, such as by decreased replication, directly or indirectly, by the TNF inhibiting-compounds of Formula (1). Such viruses include, but are not limited to HIV-1, HIV-2 and HIV-3, Cytomegalovirus (CMV), Influenza, adenovirus and the Herpes group of viruses, such as but not limited to, Herpes Zoster and Herpes Simplex. Accordingly,



in a further aspect, this invention relates to a method of treating a mammal afflicted with a human immunodeficiency virus (HIV) which comprises administering to such mammal an effective TNF inhibiting amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

5 It is also recognized that both IL-6 and IL-8 are produced during rhinovirus (HRV) infections and contribute to the pathogenesis of common cold and exacerbation of asthma associated with HRV infection (Turner *et al.*, (1998), Clin. Infec. Dis., Vol 26, p 840; Teren *et al.*, (1997), Am J Respir Crit Care Med vol 155, p1362; Grunberg *et al.* (1997), Am J Respir Crit Care Med 156:609 and Zhu *et al.*, J  
10 Clin Invest (1996), 97:421). It has also been demonstrated *in vitro* that infection of pulmonary epithelial cells with HRV results in production of IL-6 and IL-8 (Subauste *et al.*, J. Clin. Invest. 1995, 96:549.) Epithelial cells represent the primary site of infection of HRV. Therefore another aspect of the present invention is a method of treatment to reduce inflammation associated with a rhinovirus infection,  
15 not necessarily a direct effect on virus itself.

Compounds of Formula (I) may also be used in association with the veterinary treatment of mammals, other than in humans, in need of inhibition of TNF production. TNF mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted above, but in  
20 particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or retrovirus infections, such as but not limited to feline immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

25 The compounds of Formula (I) may also be used topically in the treatment or prophylaxis of topical disease states mediated by or exacerbated by excessive cytokine production, such as by IL-1 or TNF respectively, such as inflamed joints, eczema, psoriasis and other inflammatory skin conditions such as sunburn; inflammatory eye conditions including conjunctivitis; pyresis, pain and other  
30 conditions associated with inflammation. Periodontal disease has also been implemented in cytokine production, both topically and systemically. Hence use of compounds of Formula (I) to control the inflammation associated with cytokine production in such peroral diseases such as gingivitis and periodontitis is another aspect of the present invention.

35 Compounds of Formula (I) have also been shown to inhibit the production of IL-8 (Interleukin-8, NAP). Accordingly, in a further aspect, this invention relates to a method of inhibiting the production of IL-8 in a mammal in need thereof which

comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. These diseases are characterized by massive neutrophil infiltration such as, psoriasis, inflammatory bowel disease, asthma, cardiac, brain and renal reperfusion injury, adult respiratory distress syndrome, thrombosis and glomerulonephritis. All of these diseases are associated with increased IL-8 production which is responsible for the chemotaxis of neutrophils into the inflammatory site. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8 has the unique property of promoting neutrophil chemotaxis and activation. Therefore, the inhibition of IL-8 production would lead to a direct reduction in the neutrophil infiltration.

The compounds of Formula (I) are administered in an amount sufficient to inhibit cytokine, in particular IL-1, IL-6, IL-8 or TNF, production such that it is regulated down to normal levels, or in some case to subnormal levels, so as to ameliorate or prevent the disease state. Abnormal levels of IL-1, IL-6, IL-8 or TNF, for instance in the context of the present invention, constitute: (i) levels of free (not cell bound) IL-1, IL-6, IL-8 or TNF greater than or equal to 1 picogram per ml; (ii) any cell associated IL-1, IL-6, IL-8 or TNF; or (iii) the presence of IL-1, IL-6, IL-8 or TNF mRNA above basal levels in cells or tissues in which IL-1, IL-6, IL-8 or TNF, respectively, is produced.

The discovery that the compounds of Formula (I) are inhibitors of cytokines, specifically IL-1, IL-6, IL-8 and TNF is based upon the effects of the compounds of Formulas (I) on the production of the IL-1, IL-8 and TNF in *in vitro* assays which are described herein.

As used herein, the term "inhibiting the production of IL-1 (IL-6, IL-8 or TNF)" refers to:

- a) a decrease of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels by inhibition of the *in vivo* release of the cytokine by all cells, including but not limited to monocytes or macrophages;
- b) a down regulation, at the genomic level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels;
- c) a down regulation, by inhibition of the direct synthesis of the cytokine (IL-1, IL-6, IL-8 or TNF) as a posttranslational event; or
- d) a down regulation, at the translational level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels.

As used herein, the term "TNF mediated disease or disease state" refers to

any and all disease states in which TNF plays a role, either by production of TNF itself, or by TNF causing another monokine to be released, such as but not limited to IL-1, IL-6 or IL-8. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts, basophils, neutrophils, endothelial cells, brain astrocytes, bone marrow stromal cells, epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- $\alpha$ ) and Tumor Necrosis Factor beta (TNF- $\beta$ ).

As used herein, the term "cytokine interfering" or "cytokine suppressive amount" refers to an effective amount of a compound of Formula (I) which will cause a decrease in the *in vivo* levels of the cytokine to normal or sub-normal levels, when given to a patient for the prophylaxis or treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production.

As used herein, the cytokine referred to in the phrase "inhibition of a cytokine, for use in the treatment of a HIV-infected human" is a cytokine which is implicated in (a) the initiation and/or maintenance of T cell activation and/or activated T cell-mediated HIV gene expression and/or replication and/or (b) any cytokine-mediated disease associated problem such as cachexia or muscle degeneration.

As TNF- $\beta$  (also known as lymphotoxin) has close structural homology with TNF- $\alpha$  (also known as cachectin) and since each induces similar biologic responses and binds to the same cellular receptor, both TNF- $\alpha$  and TNF- $\beta$  are inhibited by the compounds of the present invention and thus are herein referred to collectively as "TNF" unless specifically delineated otherwise.

A member of the MAP kinase family, alternatively termed CSBP, p38, or RK, has been identified independently by several laboratories. Activation of this novel protein kinase via dual phosphorylation has been observed in different cell systems upon stimulation by a wide spectrum of stimuli, such as physicochemical stress and treatment with lipopolysaccharide or proinflammatory cytokines such as

interleukin-1 and tumor necrosis factor. The cytokine biosynthesis inhibitors, of the present invention, compounds of Formula (I) have been determined to be potent and selective inhibitors of CSBP/p38/RK kinase activity. These inhibitors are of aid in determining the signaling pathways involvement in inflammatory responses. In particular, for the first time a definitive signal transduction pathway can be prescribed to the action of lipopolysaccharide in cytokine production in macrophages. In addition to those diseases already noted, treatment of stroke, neurotrauma, cardiac and renal reperfusion injury, congestive heart failure, coronary arterial bypass grafting (CABG) surgery, chronic renal failure, angiogenesis & related processes, such as cancer, thrombosis, glomerulonephritis, diabetes and pancreatic  $\beta$  cells, multiple sclerosis, muscle degeneration, eczema, psoriasis, sunburn, and conjunctivitis are also included.

The CSBP inhibitors were subsequently tested in a number of animal models for anti-inflammatory activity. Model systems were chosen that were relatively insensitive to cyclooxygenase inhibitors in order to reveal the unique activities of cytokine suppressive agents. The inhibitors exhibited significant activity in many such in vivo studies. Most notable are its effectiveness in the collagen-induced arthritis model and inhibition of TNF production in the endotoxic shock model. In the latter study, the reduction in plasma level of TNF correlated with survival and protection from endotoxic shock related mortality. Also of great importance are the compounds effectiveness in inhibiting bone resorption in a rat fetal long bone organ culture system. Griswold et al., (1988) *Arthritis Rheum.* **31**:1406-1412; Badger, et al., (1989) *Circ. Shock* **27**, 51-61; Votta et al., (1994) *in vitro. Bone* **15**, 533-538; Lee et al., (1993). *B Ann. N. Y. Acad. Sci.* **696**, 149-170.

Chronic diseases which have an inappropriate angiogenic component are various ocular neovascularizations, such as diabetic retinopathy and macular degeneration. Other chronic diseases which have an excessive or increased proliferation of vasculature are tumor growth and metastasis, atherosclerosis, and certain arthritic conditions. Therefore CSBP kinase inhibitors will be of utility in the blocking of the angiogenic component of these disease states.

The term "excessive or increased proliferation of vasculature inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are characterized by hemangiomas and ocular diseases.

The term "inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are characterized by vesicle proliferation with accompanying tissue proliferation, such as occurs in cancer, metastasis, arthritis and atherosclerosis.

Another aspect of the present invention is a method of treating the common cold or respiratory viral infection caused by human rhinovirus (HRV), other enteroviruses, coronavirus, influenza virus, parainfluenza virus, respiratory syncytial virus, or adenovirus in a human in need thereof which method comprises  
5 administering to said human an effective amount of a CBSP/p38 inhibitor.

Another aspect of the present invention is a method of treating, including prophylaxis of influenza induced pneumonia in a human in need thereof which method comprises administering to said human an effective amount of a CBSP/p38 inhibitor

10 The present invention also relates to the use of the CSBP/p38 kinase inhibitor for the treatment, including prophylaxis, of inflammation associated with a viral infection of a human rhinovirus (HRV), other enteroviruses, coronavirus, influenza virus, parainfluenza virus, respiratory syncytial virus, or adenovirus.

In particular, the present invention is directed to the treatment of a viral  
15 infection in a human, which is caused by the human rhinovirus (HRV), other enterovirus, coronavirus, influenza virus, parainfluenza virus, respiratory syncytial virus, or an adenovirus. In particular the invention is directed to respiratory viral infections that exacerbate asthma (induced by such infections), chronic bronchitis, chronic obstructive pulmonary disease, otitis media, and sinusitis. While inhibiting  
20 IL-8 or other cytokines may be beneficial in treating a rhinovirus may be known, the use of an inhibitor of the p38 kinase for treating HRV or other respiratory viral infections causing the common cold is believed novel.

It should be noted that the respiratory viral infection treated herein may also be associated with a secondary bacterial infection, such as otitis media, sinusitis, or  
25 pneumonia.

For use herein treatment may include prophylaxis for use in a treatment group susceptible to such infections. It may also include reducing the symptoms of, ameliorating the symptoms of, reducing the severity of, reducing the incidence of, or any other change in the condition of the patient, which improves the therapeutic  
30 outcome.

It should be noted that the treatment herein is not directed to the elimination or treatment of the viral organism itself but is directed to treatment of the respiratory viral infection that exacerbates other diseases or symptoms of disease, such as asthma (induced by such infections), chronic bronchitis, chronic obstructive pulmonary disease, otitis  
35 media, and sinusitis.

A preferred virus for treatment herein is the human rhinovirus infection (HRV) or respiratory syncytial virus (RSV).

Another aspect of the present invention is directed to the treatment of inhaled smoke induced airway inflammation, lung chemokine production and cytokine production. The invention may be directed to treatment of the airway induced inflammation which is secondary to other respiratory disorders such as viral  
5 infections that exacerbate asthma (induced by such infections), chronic bronchitis, chronic obstructive pulmonary disease, otitis media, and sinusitis. A respiratory viral infection treated in conjunction with the smoke related airway inflammation may also be associated with a secondary bacterial infection, such as otitis media, sinusitis, or pneumonia.

10 It is noted that the inflammation may be due to cytokines and chemokine release from neutrophil activation and other leukocytes, as well as vascular and airway endothelial cell activation.

For use herein treatment may include prophylaxis for use in a treatment group who may be susceptible to such airway inflammation. It may also include  
15 reducing the symptoms of, ameliorating the symptoms of, reducing the severity of, reducing the incidence of, or any other change in the condition of the patient, which improves the therapeutic outcome.

Suitable patient populations for whom this may be prophylactically beneficial could be firemen who routinely inhale smoke in the course of their duties; use in the  
20 military, and by civilians in wartime exposure.

As noted, smoke of natural causes, such as plant extracts, natural plants products, synthetic material, chemically treated natural materials, or natural products such as oil and gas or other fossil fuels, may be treated within the scope of this invention. Suitably, the treatment including prophylaxis is related to cigarette smoke  
25 or synthetic/composites, such as occur in fires associated with burning buildings or homes.

Another aspect of the present invention relates to the use of a CSBP/p38 kinase inhibitor for the treatment, including prophylaxis, of the hypertussive activity associated with resulting airway inflammation and/or cough in a mammal in  
30 need thereof.

The present invention also relates to use of a CSBP/p38 kinase inhibitor for the treatment, including prophylaxis, of the inflammation enhanced cough related disorders in a mammal in need thereof.

The present invention is also directed to the use of a compound of Formula  
35 (I) in eosinophilic bronchitis, and in cough variant asthma.

The compounds of Formula (I) may also be used in the treatment, including prophylaxis, of eosinophilic inflammation in the airways and cough. Treatment,

including prophylaxis is appropriate for eosinophilic bronchitis (as this differs from asthma) and for the treatment, including prophylaxis of cough variant asthma.

These disorders may be directed to treatment of the airway induced inflammation which is secondary to other respiratory disorders such as viral infections that

5 exacerbate asthma (induced by such infections), chronic bronchitis, chronic obstructive pulmonary disease, otitis media, and sinusitis. A respiratory viral infection treated in conjunction with the smoke related airway inflammation may also be associated with a secondary bacterial infection, such as otitis media, sinusitis, or pneumonia.

10 The hypertussive or inflammation enhanced cough related disorders may either be a direct result of or an association with eosinophilia activity. It may also be a result of, or associated with the blocking production of certain cytokines which may mediate these phenomena.

For use herein treatment may include prophylaxis for use in a treatment  
15 group who may be susceptible to such airway inflammation, and/or cough. It may also include reducing the symptoms of, ameliorating the symptoms of, reducing the severity of, reducing the incidence of, or any other change in the condition of the patient, which improves the therapeutic outcome.

Clinically, eosinophilic bronchitis presents as chronic cough and sputum  
20 eosinophilia, but without the abnormalities of airway function seen in asthma. In contrast to cough in patients without sputum eosinophilia, the cough responds to anti-inflammatory therapy, such as inhaled corticosteroids (Niimi et al., Eosinophilic inflammation in cough variant asthma, European Respiratory Journal. 11(5):1064-9, (1998)).

25 Patients with cough-variant asthma may also have the following criteria: (1) have not been previously diagnosed as having asthma; (2) complain of a cough of at least a 3-week duration; (3) do not complain of wheezing, shortness of breath, or chest tightness; (4) have normal results of physical examinations; (5) have normal or nearly normal results of spirometry; (6) have evidence of bronchial  
30 hyperresponsiveness during bronchoprovocation challenge testing; and (7) have a favorable response to asthma medications (Irwin et al., Interpretation of positive results of a methacholine inhalation challenge and 1 week of inhaled bronchodilator use in diagnosing and treating cough-variant asthma (Archives of Internal Medicine. 157(17):1981-1987, (1997)).

35 Unlike conventional anti-tussive agents, such as codeine or dextromethorphan, a p38 kinase inhibitor appears to have no direct antitussive activity, but reduces the airway eosinophilia and normalizes the hypertussive state. Therefore, use of a p38

inhibitor will reduce the added coughs, or hypertussive state, back to a normal level which can be suitably treated with conventional agents and/or therapies as appropriate. Use of the p38 inhibitors will allow for the maintenance of patients who are subject to increased cough responsiveness, especially unproductive cough, due to other underlying disorders or treatments. This increased cough responsiveness may be modulated, or  
5 decreased by use of this innovative anti-inflammatory therapy.

Accordingly, the present invention provides a method of treating a CSBP kinase mediated disease in a mammal in need thereof, preferably a human, which comprises administering to said mammal, an effective amount of a compound of  
10 Formula (I) or a pharmaceutically acceptable salt thereof.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. This invention, therefore, also relates to a pharmaceutical composition comprising an effective, non-  
15 toxic amount of a compound of Formula (I) and a pharmaceutically acceptable carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally,  
20 topically, parenterally or by inhalation. The compounds of Formula (I) may be administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active  
25 compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The carrier(s)  
30 must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid  
35 carriers are syrup, peanut oil, olive oil, H<sub>2</sub>O and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.



A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg. to about 1g. When  
5 a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I)  
10 externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid  
15 preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the formulation. It may however comprise as much as 10% w/w but  
20 preferably will comprise less than 5% w/w, more preferably from 0.1% to 1% w/w of the formulation.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those  
25 for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid  
30 formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage;  
35 an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or a macrogel. The formulation may incorporate any suitable

surface active agent such as an anionic, cationic or non-ionic surfactant such as a sorbitan ester or a polyoxyethylene derivative thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

5           Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a  
10   suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C. for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and  
15   chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.

          Compounds of Formula (I) may be administered parenterally, that is by intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of  
20   parenteral administration are generally preferred. Appropriate dosage forms for such administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation, that is by intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional  
25   techniques.

          For all methods of use disclosed herein for the compounds of Formula (I), the daily oral dosage regimen will preferably be from about 0.1 to about 80 mg/kg of total body weight, preferably from about 0.2 to 30 mg/kg, more preferably from about 0.5 mg to 15mg. The daily parenteral dosage regimen about 0.1 to about 80  
30   mg/kg of total body weight, preferably from about 0.2 to about 30 mg/kg, and more preferably from about 0.5 mg to 15mg/kg. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be recognized by one of skill in  
35   the art that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of the condition being treated, the form, route and site of

administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The novel compounds of Formula (I) may also be used in association with the veterinary treatment of mammals, other than humans, in need of inhibition of CSBP/p38 or cytokine inhibition or production. In particular, CSBP/p38 mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section, but in particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or retrovirus infections, such as but not limited to feline immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

#### BIOLOGICAL EXAMPLES

The cytokine-inhibiting effects of compounds of the present invention may be determined by the following *in vitro* assays:

Assays for Interleukin - 1 (IL-1), Interleukin -8 (IL-8 ), and Tumour Necrosis Factor (TNF) are well known in the art, and may be found in a number of publications, and patents. Representative suitable assays for use herein are described in Adams et al., US 5,593,992, whose disclosure is incorporated by reference in its entirety.

##### **Interleukin - 1 (IL-1)**

Human peripheral blood monocytes are isolated and purified from either fresh blood preparations from volunteer donors, or from blood bank buffy coats, according to the procedure of Colotta *et al*, J Immunol, **132**, 936 (1984). These monocytes ( $1 \times 10^6$ ) are plated in 24-well plates at a concentration of 1-2 million/ml per well. The cells are allowed to adhere for 2 hours, after which time non-adherent cells are removed by gentle washing. Test compounds are then added to the cells for 1h before the addition of lipopolysaccharide (50 ng/ml), and the cultures are incubated at 37°C for an additional 24h. At the end of this period, culture supernatants are removed and clarified of cells and all debris. Culture supernatants are then immediately assayed for IL-1 biological activity, either by the method of Simon *et al.*, J. Immunol. Methods, **84**, 85, (1985) (based on ability of IL-1 to

stimulate a Interleukin 2 producing cell line (EL-4) to secrete IL-2, in concert with A23187 ionophore) or the method of Lee *et al.*, J. ImmunoTherapy, **6** (1), 1-12 (1990) (ELISA assay).

**In vivo TNF assay:**

5 (1) Griswold *et al.*, Drugs Under Exp. and Clinical Res., **XIX** (6), 243-248 (1993); or

(2) Boehm, *et al.*, *Journal Of Medicinal Chemistry* **39**, 3929-3937 (1996) whose disclosures are incorporated by reference herein in their entirety.

**LPS-induced TNF $\alpha$  Production in Mice and Rats**

10 In order to evaluate in vivo inhibition of LPS-induced TNF $\alpha$  production in rodents, both mice and rats are injected with LPS.

Mouse Method

Male Balb/c mice from Charles River Laboratories are pretreated (30 min) with compound or vehicle. After the 30 min. pretreat time, the mice are given LPS (lipopolysaccharide from Esherichia coli Serotype 055-85, Sigma Chemical Co., St Louis, MO) 25 ug/mouse in 25 ul phosphate buffered saline (pH 7.0) intraperitoneally. Two hours later the mice are killed by CO<sub>2</sub> inhalation and blood samples are collected by exsanguination into heparinized blood collection tubes and stored on ice. The blood samples are centrifuged and the plasma collected and stored at -20°C until assayed for TNF $\alpha$  by ELISA.

Rat Method

Male Lewis rats from Charles River Laboratories are pretreated at various times with compound or vehicle. After a determined pretreat time, the rats are given LPS (lipopolysaccharide from Esherichia coli Serotype 055-85, Sigma Chemical Co., St Louis, MO) 3.0 mg/kg intraperitoneally. The rats are killed by CO<sub>2</sub> inhalation and heparinized whole blood is collected from each rat by cardiac puncture 90 min after the LPS injection. The blood samples are centrifuged and the plasma collected for analysis by ELISA for TNF $\alpha$  levels.

ELISA Method

30 TNF $\alpha$  levels were measured using a sandwich ELISA, as described in Olivera *et al.*, *Circ. Shock*, **37**, 301-306, (1992), whose disclosure is incorporated by reference in its entirety herein, using a hamster monoclonal antimurine TNF $\alpha$  (Genzyme, Boston, MA) as the capture antibody and a polyclonal rabbit antimurine TNF $\alpha$  (Genzyme) as the second antibody. For detection, a peroxidase-conjugated goat antirabbit antibody (Pierce, Rockford, IL) was added, followed by a substrate for peroxidase (1 mg/ml orthophenylenediamine with 1% urea peroxide). TNF $\alpha$  levels in the plasma samples from each animal were calculated from a standard curve generated

with recombinant murine TNF $\alpha$  (Genzyme).

**LPS-Stimulated Cytokine Production in Human Whole Blood**

Assay: Test compound concentrations were prepared at 10 X concentrations and LPS prepared at 1 ug/ml (final conc. of 50 ng/ml LPS) and added in 50 uL volumes to 1.5 mL eppendorf tubes. Heparinized human whole blood was obtained from healthy volunteers and was dispensed into eppendorf tubes containing compounds and LPS in 0.4 mL volumes and the tubes incubated at 37 C. Following a 4 hour incubation, the tubes were centrifuged at 5000 rpm for 5 min in a TOMY microfuge, plasma was withdrawn and frozen at -80 C.

Cytokine measurement: IL-1 and/or TNF were quantified using a standardized ELISA technology. An in-house ELISA kit was used to detect human IL-1 and TNF. Concentrations of IL-1 or TNF were determined from standard curves of the appropriate cytokine and IC<sub>50</sub> values for test compound (concentration that inhibited 50% of LPS-stimulated cytokine production) were calculated by linear regression analysis.

**CSBP/p38 Kinase Assay:**

This assay measures the CSBP/p38-catalyzed transfer of <sup>32</sup>P from [ $\alpha$ -<sup>32</sup>P]ATP to threonine residue in an epidermal growth factor receptor (EGFR)-derived peptide (T669) with the following sequence: KRELVEPLTPSGEAPNQALLR (residues 661-681). (See Gallagher *et al.*, "Regulation of Stress Induced Cytokine Production by Pyridinyl Imidazoles: Inhibition of CSBP Kinase", BioOrganic & Medicinal Chemistry, 1997, 5, 49-64).

Reactions were carried in round bottom 96 well plate (from Corning) in a 30 ml volume. Reactions contained (in final concentration): 25 mM Hepes, pH 7.5; 8 mM MgCl<sub>2</sub>; 0.17 mM ATP (the K<sub>m</sub>[ATP] of p38 (see Lee et al., Nature 300, n72 pg. 639-746 (Dec. 1994))); 2.5 uCi of [ $\gamma$ -<sup>32</sup>P]ATP; 0.2 mM sodium orthovanadate; 1 mM DTT; 0.1% BSA; 10% glycerol; 0.67 mM T669 peptide; and 2-4 nM of yeast-expressed, activated and purified p38. Reactions were initiated by the addition of [ $\gamma$ -<sup>32</sup>P]Mg/ATP, and incubated for 25 min. at 37 °C. Inhibitors (dissolved in DMSO) were incubated with the reaction mixture on ice for 30 min prior to adding the <sup>32</sup>P-ATP. Final DMSO concentration was 0.16%. Reactions were terminated by adding 10 ul of 0.3 M phosphoric acid, and phosphorylated peptide was isolated from the reactions by capturing it on p81 phosphocellulose filters. Filters were washed with 75 mM phosphoric acids, and incorporated <sup>32</sup>P was quantified using beta scintillation counter. Under these conditions, the specific activity of p38 was 400-450 pmol/pmol enzyme, and the activity was linear for up to 2 hours of incubation. The kinase activity values were obtained after subtracting values

generated in the absence of substrate which were 10-15% of total values.

Representative compounds of Formula (I), Examples 1 to 167 have demonstrated positive inhibitory activity in this assay, all having of IC<sub>50</sub>'s of < 100uM in this binding assay.

#### 5 **TNF- $\alpha$ in Traumatic Brain Injury Assay**

This assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions which follow experimentally induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Since TNF- $\alpha$  is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated  
10 astrocytes, this post-traumatic alteration in gene expression of TNF- $\alpha$  plays an important role in both the acute and regenerative response to CNS trauma. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

#### **CNS Injury model for IL-b mRNA**

15 This assay characterizes the regional expression of interleukin-1 $\beta$  (IL-1 $\beta$ ) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Results from these assays indicate that following TBI, the temporal expression of IL-1 $\beta$  mRNA is regionally stimulated in specific brain regions. These regional changes in cytokines, such as IL-1 $\beta$  play a role in the post-  
20 traumatic pathologic or regenerative sequelae of brain injury. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

#### **Angiogenesis Assay**

Described in WO 97/32583, whose disclosure is incorporated herein by reference, is an assay for determination of inflammatory angiogenesis which may be  
25 used to show that cytokine inhibition will stop the tissue destruction of excessive or inappropriate proliferation of blood vessels.

#### **Cigarette Smoke Exposure Model**

A murine model of cigarette smoke inhalation was developed to explore a relationship to leukocyte trafficking and lung chemokine and cytokine production.  
30 Balb/c mice are exposed to smoke generated from commercial unfiltered cigarettes for a specified period of time and samples are obtained at varying times during the post-exposure. This model is demonstrated in greater detail as shown below, in contrast to other smoke extract models known in the art.

A model of cigarette smoke exposure in the mouse is established in which  
35 mice are placed 6 at a time into a small animal plexiglass dosing chamber attached to a peristaltic pump whose intake is connected to a holder for a commercial unfiltered cigarette (Lucky Strike<sup>TM</sup>). Along with fresh air, smoke is delivered into

the chamber until the cigarette is consumed (approximately 5 minutes). Varying numbers of cigarettes (2-4 per day, 2-3 hr apart) are utilized for 1-3 consecutive days. Animals are euthanized by pentobarbital overdose approximately 18 hours after the final exposure. Bronchoalveolar lavage with phosphate-buffered saline is performed for inflammatory cell enumeration, and BAL aliquots and lungs are frozen for cytokine analysis. Smoke exposure results in time- and cigarette number-related increases in airway neutrophils, and lung chemokine (KC) and cytokine (IL-6) content.

To evaluate the role of a p38 MAP kinase inhibitor in this inflammatory response, mice are treated with a p38 kinase inhibitor, a compound of Formula (I) at approximately a 30 mg/kg, p.o. b.i.d.. Reduction in lung KC (a murine homologue of IL-8) levels are assessed 1 day after exposure (prior to neutrophilia), and attenuated airway neutrophilia and lung IL-6 levels are assessed following 3 days of cigarette exposure.

#### **Hypertussive Cough Models**

Described below is an example of how to determine the usefulness of p38 inhibitors in the treatment of hypertussive disorders or inflammation enhanced cough.

The directed antitussive activity of the compound in question is first assessed, by a 10 to 30 minute pretreatment period by intraperitoneal injection or a 1 hour pretreatment period for oral administration. The animals (guinea pigs) are then subjected to an inhaled citric acid-induced cough challenge. The Citric Acid Induced Cough Model is shown in Figure 2.

The effects of the compound are then assessed on the hypertussive response that occurs 72 hours post aerosol exposure to antigen or LTD4 exposure. Treatment of the animals occurs with the drug prior and/or after antigen or LTD4 challenge, but not on the day of citric acid challenge. The antigen or LTD4 induced hypertussive model is shown in Figure 3.

The effects of known antitussive agents, dextromethorphan and codeine on Citric Acid Induced Cough in Guinea Pigs is shown in Figure 4.

Inhalation of citric acid (CA; 0.4% for 1 minute) induces 11 to 15 coughs during the exposure and 12 -minute monitoring period in conscious guinea pigs. Exposure of sensitized animals to inhaled ovalbumin resulted in a hypertussive state (50-80% increase in CA-induced cough incidence) for several days, which positively correlated with airway eosinophilia determined by bronchoalveolar lavage.

Similarly, inhalation of LTD4 (10 ug/ml for 1 minute) increases cough incidence and airway eosinophils 72 hours after exposure. Further explanation and details may be found in PCT/US00/25386, filed 15 September 2000 whose

disclosure is incorporated by reference in its entirety.

## SYNTHETIC EXAMPLES

5 The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. All temperatures are given in degrees centigrade, all solvents are highest available purity and all reactions run under anhydrous conditions in an argon atmosphere where necessary.

Mass spectra were run on an open access LC-MS system using electrospray  
10 ionization. LC conditions: 4.5% to 90% CH<sub>3</sub>CN (0.02% TFA) in 3.2 min with a 0.4 min hold and 1.4 min re-equilibration; detection by MS, UV at 214, and a light scattering detector (ELS). Column: 1 X 40 mm Aquasil (C18) <sup>1</sup>H-NMR (hereinafter "NMR") spectra were recorded at 400 MHz using a Bruker AM 400 spectrometer or a Bruker AVANCE 400. Multiplicities indicated are: s=singlet, d=doublet, t=triplet, q=quartet,  
15 m=multiplet and br indicates a broad signal. Preparative (prep) hplc; ca 50 mg of the final products were injected in 500 uL of DMSO onto a 50 X 20 mm I. D. YMC CombiPrep ODS-A column at 20 mL/min with a 10 min gradient from 10% CH<sub>3</sub>CN (0.1% TFA) to 90% CH<sub>3</sub>CN (0.1% TFA) in H<sub>2</sub>O (0.1% TFA) and a 2 min hold. Flash chromatography is run over Merck Silica gel 60 (230 - 400 mesh).

20 Accepted abbreviations are used as described in the ACS Style Guide (*The ACS Style Guide. A Manual for Authors*; Dodd, J. A., Ed.; American Chemical Society: Washington, DC, 1986; pp 47-69.). In addition the following are used:

BOC t-butoxycarbonyl  
eq indicates the proportion of molar equivalents of a reagent relative to the principal  
25 reagent  
NMP 1-methyl-2-pyrrolidinone  
satd saturated  
Rt hplc retention time

### Example 1

30 Preparation of 1,5-diphenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

a) 4-Chloro-2-methylsulfanyl-6-phenylamino-pyrimidine-5-carbonitrile  
4,6-Dichloro-2-methylsulfanyl-pyrimidine-5-carbonitrile [Santilli, et al., *J. Heterocycl. Chem.* **1971**, 8, 445-453] (0.222 grams (hereinafter "g"), 1.0 millimoles  
35 (hereinafter "mmol")) in EtOH (2 milliliters (hereinafter "mL")) and Et<sub>2</sub>O (1mL) was treated with aniline (184 microliter (hereinafter "uL"), 2.0 mmol) in EtOH (1 mL). A clear solution formed initially but rapidly formed a heavy precipitate. The mixture was



stirred 30 minutes (hereinafter "min") and was filtered, and the solid was washed with 1:1 Et<sub>2</sub>O, EtOH and then Et<sub>2</sub>O, dried to afford 162 milligrams (hereinafter "mg") (59%) of the title compound as a white solid. LC MS m/e = 277 (MH<sup>+</sup>), Rt = 2.32 min.

b) 2-Methylsulfanyl-4-phenyl-6-phenylamino-pyrimidine-5-carbonitrile

5 The product of the previous example, (162 mg, 0.59 mmol), phenylboronic acid (360 mg, 2.95 mmol), Na<sub>2</sub>CO<sub>3</sub> (318 mg, 3 mmol), dioxane (3 mL) and H<sub>2</sub>O (1.5 mL) were combined and argon was bubbled through the mixture for 30 min Pd[P(Ph)<sub>3</sub>] (15 mg, 0.013 mmol) was added and the mixture was heated to 85° for 1.5 h, then cooled, diluted with H<sub>2</sub>O (25 mL) and extracted with EtOAc (6 X 50 mL). The combined  
10 extracts were washed with H<sub>2</sub>O, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and filtered through a 10 g plug of silica (Varien bond elute<sup>®</sup>) with CH<sub>2</sub>Cl<sub>2</sub> to afford the title compound as a white solid 185 mg (97%). LC MS m/e = 319 (MH<sup>+</sup>), Rt = 2.59 min.

c) (5-Aminomethyl-2-methylsulfanyl-6-phenyl-pyrimidin-4-yl)-phenylamine.

The product of the previous example (151 mg, 0.47 mmol) was dissolved in warm  
15 dioxane (6 mL), cooled to 23° and 1M LAH in Et<sub>2</sub>O (1 mL, 1 mmol) was added and the resulting solution was heated to 55° for 1.5 h., diluted with EtOAc (10 mL) and then poured into 10% aq NaOH (20 mL) and extracted with more EtOAc (2 X 50 mL). The combined EtOAc was washed with H<sub>2</sub>O, then satd aq NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford the title compound as a yellow solid. 153 mg (100%). LC MS m/e  
20 = 323 (MH<sup>+</sup>), Rt = 1.49 min

d) 1,5-diphenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of the previous example (153 mg, 0.47 mmol) was dissolved in toluene (5 mL) and pyridine (2 mL). A 20% solution of COCl<sub>2</sub> in toluene (2 mL) was added and the resulting mixture was stirred 30 min, diluted with EtOAc (125 mL) and  
25 washed with 10% aq NaOH (20 mL) and H<sub>2</sub>O (5 X 20 mL) and concentrated. The crude product was purified by chromatotron chromatography (0 - 2 % MeOH in CH<sub>2</sub>Cl<sub>2</sub>) and then the resulting nearly pure yellow solid was further purified by prep hplc. The resulting white foam was dried *in vacuo*, triturated with H<sub>2</sub>O, and filtered and dried *in vacuo* to afford 33 mg of the title compound as a white powder. LC MS m/e = 349  
30 (MH<sup>+</sup>), Rt = 2.12 min.

#### Example 2

1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

a) 4-Chloro-6-(2,6-difluoro-phenylamino)-2-methylsulfanyl-pyrimidine-5-carbonitrile.

2,6-di-Fluoroaniline (2.66 mL, 24 mmol), DMSO (12 mL) and NaH (0.912 g, 22.8 mmol) were combined at 23°. When the foaming ceased 4,6-Dichloro-2-

methysulfanyl-pyrimidine-5-carbonitrile [Santilli, et al., *J. Heterocycl. Chem.* **1971**, 8, 445-453] (5.0 g, 22.82 mmol) was added in DMSO (12 mL). The reaction exothermed and the temperature was moderated with cold H<sub>2</sub>O bath, then stirred at 23° for 3 h. The reaction was diluted with EtOAc (300 mL) and washed with H<sub>2</sub>O (4X) and satd aq NaCl (1X). Dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford 6.21 g (82%) of a tan foam. LC MS m/z = 313 (MH<sup>+</sup>), Rt = 2.37 min.

b) 4-(2,6-Difluorophenylamino)-6-(4-fluoro-2-methylphenyl)-2-methysulfanylpurimidine-5-carbonitrile.

The product of the previous example (4.0 g, 12.8 mmol), 2-methyl-4-fluorophenylboronic acid (Lancaster) (5.91 g, 38.4 mmol), Na<sub>2</sub>CO<sub>3</sub> (4.04 g, 38.4 mmol), dioxane (100 mL) and H<sub>2</sub>O (50 mL) were combined and a stream of argon was passed through the mixture for 15 min. Pd[P(Ph)<sub>3</sub>]<sub>4</sub> (Aldrich) (400 mg) was added and the mixture was heated to 85° for 6 h. EtOAc (600 mL) was added and the organic phase was washed with H<sub>2</sub>O (100 mL). The aq wash was extracted with EtOAc (2 X 75 mL) and the combined organic phases were washed with more H<sub>2</sub>O (50 mL) and satd aq NaCl (50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and the residue was flash chromatographed with CH<sub>2</sub>Cl<sub>2</sub>. The desired fractions were pooled and concentrated *in vacuo* to afford 4.16 g (84%) of a tan foam. LC MS m/z = 387 (MH<sup>+</sup>), Rt = 2.52 min

c) 5-Aminomethyl-6-(4-fluoro-2-methylphenyl)-2-methysulfanylpurimidin-4-yl]-(2,6-difluorophenyl)amine.

The product of the previous example (4.24 g, 11.0 mmol) was dissolved in Et<sub>2</sub>O (200 mL) and 1M LAH in Et<sub>2</sub>O (22 mL) was added dropwise. After addition the mixture was heated to Et<sub>2</sub>O reflux for 1.5 h, cooled to 4° and quenched by the addition of H<sub>2</sub>O (1.1 mL), then 10% aq NaOH (5.5 mL) and then more H<sub>2</sub>O (5.5 mL), stirred 10 min then 2.5% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub> (400 mL) was added. Stirred 15 min and the organic layer was decanted and the residue was washed with additional 2.5% methanolic CH<sub>2</sub>Cl<sub>2</sub> (200 mL). The combined organic layers were dried (Na<sub>2</sub>SO<sub>2</sub>) and concentrated to afford 4.02 g (94%) of a yellow foamy solid. LC MS m/z = 391 (MH<sup>+</sup>), Rt = 1.62 min.

d) 1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methysulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of the previous example, (4.02 g, 10.4 mmol) was dissolved in THF (100 mL) and carbonyldiimidazole (2.23 g, 13.8 mmol) was added. The resulting mixture was stirred 16h, diluted with EtOAc (1 L) and washed with H<sub>2</sub>O (4 X 150 mL) and satd aq NaCl (150 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to a brown foam. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and flash chromatographed in 0 - 2% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub> to afford 2.97 g (69%) of the title compound as an off white solid. LC MS m/z = 417 (MH<sup>+</sup>), Rt = 2.27 min.

Example 31-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of the preceding example, example 2 (2.14 g, 5.14 mmol), CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and *meta*-chloroperoxybenzoic acid (57 - 85%) (3.102 g, 10.26 mmol if 57%) were dissolved together. After 16 h, the reaction was diluted with EtOAc (300 mL) and washed with 5% aq Na<sub>2</sub>CO<sub>3</sub> (6 X 50 mL), H<sub>2</sub>O (50 mL) and satd aq NaCl (50 mL), dried Na<sub>2</sub>SO<sub>2</sub> and concentrated to afford 2.3 g (100 %) of the title compound as a white solid. LC MS m/z = 417 (MH<sup>+</sup>), Rt = 1.94 min

Example 41-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(piperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

a) 1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-*tert*-butoxycarbonyl-piperidin-4-ylamino)-3,4-dihydro-pyrimido[4,5-d]pyrimidin-2-one

The product of the previous example, Example 3 (90 mg, 0.2 mmol), NMP (1 mL) and 4-amino-1-BOC-piperidine (Astatech) (200 mg, 1.0 mmol) were heated in an oil bath at 65° for 16 h. The reaction was cooled to 23°, diluted with EtOAc (75 mL) and washed with 10 % aq citric acid (2X), H<sub>2</sub>O and satd aq NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to afford 112 mg of a white solid. LC MS m/z = 569 (MH<sup>+</sup>), Rt = 2.14 min

b) 1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(piperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

The product of the preceding example was dissolved in TFA (5 mL) and allowed to stand for 15 min, then concentrated and purified by prep hplc to afford 56.8 mg (61%) of a white powder. LC MS m/z = 469 (MH<sup>+</sup>), Rt = 1.35 min

Example 51-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-dimethylaminoethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

The product of example 3 (90 mg, 0.2 mmol) was dissolved in NMP (1 mL) and dimethylaminoethylamine (110 uL, 1.0 mmol) was added and the resulting brown soln was stirred on an oil bath heated to 65° for 16 h., diluted with EtOAc (75 mL) and washed with H<sub>2</sub>O (3X) and satd aq NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>) concentrated and purified by prep hplc to afford 68 mg (75%). LC MS m/z = 457 (MH<sup>+</sup>), Rt = 1.39 min

Example 61-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1,3-dihydroxyprop-2-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of example 3 (90 mg, 0.2 mmol) was dissolved in NMP (1 mL) and serinol (91 mg, 1.0 mmol) was added. The solution was heated to 65° for 48 h, diluted

with EtOAc (75 mL) and washed with 10% aq citric acid (2X), H<sub>2</sub>O, and satd aq NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and purified by prep hplc to afford 46 mg (50%). LC MS m/z = 460 (MH<sup>+</sup>), Rt = 1.37 min

The compounds in Table 1 were prepared by either the method of Example 5 (method A) or by the method of Example 6 (method B), using the appropriate amine. Note that Methods A and B differ by the absence (method A) or presence (method B) of an aqueous citric acid wash during work-up. For those examples where the reaction time and/or temperature was varied from those given in Examples 5 and 6, the changes are indicated in Table 1, below.

10

**Table 1**

Example	Compound name	Method (variations)	LC MS	
			m/z	Rt min
7	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperazin-4-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	469	1.49
8	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(imidazol-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A (5 h)	437	1.64
9	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-([2-hydroxyethyl]-methylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (2 h)	444	1.65
10	1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-hydroxypiperdin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (4 h)	470	1.79
11	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-carboethoxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (1 h)	526	2.42
12	1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-hydroxymethylpiperdin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (1 h)	484	2.49
13	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-di(ethan-2-ol-)amino-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (24 H)	474	1.55

Table 1 cont.

Example	Compound name	Method (variations)	LC MS	
			m/z	Rt min
14	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-yl-amino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	483	1.37
15	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-morpholin-4-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (16 h)	456	2.12
16	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-morpholin-4-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	499	1.40
17	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	483	1.50
18	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	497	1.55
19	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyridin-3-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	491	1.47
20	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-diethylamino ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	485	1.55
21	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyridin-3-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A	491	1.47
22	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-dimethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A <sup>1</sup>	414	1.97
23	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-methylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	A <sup>2</sup>	400	1.64

Table 1 cont.

Example	Compound name	Method (variations)	LC MS	
			m/z	Rt min
24	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-methyl-1,3-dihydroxyprop-2-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (80°, 24h)	474	1.45
25	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (2 h)	444	1.55
26	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (2 h)	444	1.55
27	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (80°, 4 h)	458	1.57
28	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (85°, 16 h)	490	1.19
29	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (0.5 h)	430	1.50
30	1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2,3-dihydroxypropylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	B (16 h)	460	1.32

1. Using 40% aq dimethylamine. 2. Using 2M methylamine in THF.

### Example 31

#### 5 1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfinyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of example 2 (204 mg, 0.49 mmol) in THF was treated dropwise with oxone® (0.307 g, 0.5 mmol) in H<sub>2</sub>O (10 mL). The mixture was stirred at 23° for 2 h, diluted with EtOAc (100 mL) and washed with H<sub>2</sub>O (2 x 25 mL) and satd aq NaCl (25 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated. The residue was purified by prep hplc to afford the sulfoxide as a white powder. LC MS m/z = 433 (MH<sup>+</sup>), Rt = 1.70 min

10

Example 32

1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methyl-1H-imidazol-2-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

N-Methylimidazole (82  $\mu$ L, 1.0 mmol) in dry THF was cooled to  $-70^{\circ}$  and n-butyl lithium (2.5 M in hexane) (0.36 mL, 0.9 mmol) was added, the mixture was stirred 15 min and then the product of example 3 (45 mg, 0.1 mmol) was added and the reaction was warmed to  $23^{\circ}$ , and stirred 30 min. The reaction was poured into satd aq  $\text{NaHCO}_3$  (20 mL) and stirred 5 min and then extracted with EtOAc (2 X). The combined EtOAc was washed with  $\text{H}_2\text{O}$  (3X), satd aq  $\text{NaCl}$ , dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was purified by prep hplc to afford 11.6 mg (26%) of the title compound as a white powder. LC MS  $m/z$  = 451 ( $\text{MH}^+$ ),  $R_t$  = 1.55 min.

Example 33

1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-piperazin-4-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The title compound was prepared by the method of example 4 except that 1-BOC-piperazine (Fluka) was the amine. TFA deblocking of the intermediate BOC protected compound as in example 4 and prep hplc afforded the the title compound as a white powder. LC MS  $m/z$  = 455 ( $\text{MH}^+$ ),  $R_t$  = 1.55 min.




Example 34

1-(2,6-difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-carboxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of example 11 (56 mg, 0.11 mmol) was dissolved in THF (2 mL) and LiOH (24 mg, 1.0 mmol) in  $\text{H}_2\text{O}$  (1.0 mL) was added, and the resulting soln was stirred for 4 h. The reaction was concentrated and redissolved in DMSO and purified by prep hplc to afford 24.8 mg (45%) of the title compound. LC MS  $m/z$  = 498 ( $\text{MH}^+$ ),  $R_t$  = 1.64 min.

The sulfides in examples 35 to 37 (**Table 2**) were prepared by the method of Example 2, except that the boronic acid in step 2b was varied as depicted in **Table 2**. below.

**Table 2**

Example	Compound name	Boronic acid	LC MS	
			m/z	Rt (min)
35	1-(2,6-Difluorophenyl)- 5-phenyl-7-methyl-sulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one		385	2.19
36	1-(2,6-Difluorophenyl)- 5-(4-fluorophenyl)- 7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one		403	2.19
37	1-(2,6-difluorophenyl)-5-(2-methylphenyl)-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one		399	2.20

The sulfides in examples 38 - 44 (**Table 3**) were prepared by the method of Example 2 except that the amine in step 2a was varied as depicted in **Table 3**, below.

**Table 3**

Example	Compound Name	Amine	LC MS	
			m/z	Rt (min)
38	1-((R)-1-Phenylethyl -5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	R- $\alpha$ -methyl-benzylamine	409	2.39
39	1-((S)-1-Phenylethyl -5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	S- $\alpha$ -methyl-benzylamine	409	2.39
40	1-(2-Chlorophenyl)- 5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	2-chloroaniline	415	2.17



Table 3 cont.

Example	Compound Name	Amine	LC MS	
			m/z	Rt (min)
41	1-Cyclohexyl -5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	cyclohexyl-amine	387	2.62
42	1-(2-Methylphenyl)- 5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	2-methylaniline	395	2.20
43	1-(2,6-Dimethylphenyl)- 5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	2,6-dimethylaniline	409	2.22
44	1-(2-Fluorophenyl)- 5-(2-methyl-4-fluoro)phenyl-7-methylsulfanyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	2-fluoroaniline	399	2.10

The sulfones in **Table 4** were made by the procedure of Example 3 except that the substrate sulfides were the products of the examples indicated.

5

**Table 4**

Example	Compound name	Method*	LC MS	
			m/z	Rt (min)
45	1-(2,6-Difluorophenyl)- 5-phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	35	417	1.75
46	1-(2,6-Difluorophenyl)- 5-(4-fluorophenyl)- 7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	36	436	1.82

Table 4 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt (min)
47	1-(2,6-Difluorophenyl)-5-(2-methylphenyl)-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	37	431	1.87
48	1-((R))-1-Phenylethyl-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	38	441	2.01
49	1-((S))-1-Phenylethyl-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	39	441	2.01
50	1-(2-Chlorophenyl)-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	40	447	1.94
51	1-Cyclohexyl-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	41	419	2.02
52	1-(2-Methylphenyl)-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	42	427	1.82
53	1-(2,6-Dimethylphenyl)-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	43	441	1.92
54	1-(2-Fluorophenyl)-5-(2-methyl-4-fluoro)phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	44	431	1.84

\* the substrate is the product of the noted example #

#### Example 55

5 1-(2,6-Difluorophenyl)-5-phenyl-7-(4-methylpiperazin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of example 45 (41.6 mg, 0.1 mmol), 4-methylpiperazine (30 mg, 0.3 mmol) and NMP (1.0 mL) were dissolved together and heated to 65° for 18 h. The NMP was removed in high vacuum and the residue was purified by prep hplc to afford 17 mg of the title compound. LC MS m/z = 437 (MH<sup>+</sup>), Rt = 1.47 min.

Example 561-(2,6-Difluorophenyl)-5-phenyl-7-piperazin-1-yl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of example 45 (41.6 mg, 0.1 mmol), and 1-BOC-piperazine (Fluka) were treated by the method of example 55, omitting the hplc step, to afford the BOC protected intermediate. This product was deblocked by the procedure of example 4b and purified by prep hplc to afford the title compound. LC MS  $m/z$  = 422 ( $MH^+$ ),  $R_t$  = 1.37 min.

The compounds in **Table 5** were prepared by either the method of Example 5, using the appropriate amine and the sulfone shown in **Table 5** (Method C; includes aqueous work-up), or the method of Example 55 using the appropriate amine and the sulfone shown in **Table 5** (Method D; with concentration of the crude reaction mixture, then preparation hplc), or the method of Example 33 using the appropriate sulfone (Method E; aq. work-up with citric acid, deblocking, and prep hplc) or the method of Example 56 using the appropriate sulfone as shown in **Table 5** (Method F; concentration, deblocking, and prep hplc). In the table, method\* is the the substrate is the sulfone of the noted example.

**Table 5**

Example	Compound name	Method*	LC MS	
			$m/z$	$R_t$
57	1-(2,6-difluorophenyl)-5-(2-methylphenyl)-7-(2-diethylaminoethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate	C (47)	467	1.44
58	1-(2,6-difluorophenyl)-5-(2-methylphenyl)-7-(1-piperazin-4-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	E (47)	437	1.50
59	1-(2,6-difluorophenyl)-5-phenyl-7-(2-hydroxyethyl)methylamino-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (45)	412	1.60
60	1-(2,6-difluorophenyl)-5-phenyl-7-(4-carboethoxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (45)	494	1.85

Table 5 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt
61	1-(2,6-difluorophenyl)-5-phenyl-7-(4-hydroxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (45)	438	1.74
62	1-(2,6-difluorophenyl)-5-phenyl-7-(imidazol-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (45)	405	1.47
63	1-(2,6-difluorophenyl)-5-phenyl-7-(morpholin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (45)	424	2.04
64	1-(2,6-Difluorophenyl)-5-(4-fluorophenyl)-7-(4-methylpiperazin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	455	1.47
65	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(2-hydroxyethyl)methylamino-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	430	1.72
66	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(piperazin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	F (46)	441	1.40
67	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(4-carboethoxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	512	2.42
68	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(4-hydroxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	456	1.95
69	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(imidazol-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	423	1.52
70	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(morpholin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	442	2.10
71	1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(4-hydroxymethylpiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one	D (46)	470	1.90

Example 72

1-(2,6-difluorophenyl)-5-phenyl-7-(1-methyl-1H-imidazol-2-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

5 The product of example 45 was converted to the target compound by the procedure of example 32. LC MS m/z = 419 (MH<sup>+</sup>), Rt = 1.42 min

Example 73

1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(1-methyl-1H-imidazol-2-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

10 The product of example 46 was converted to the target compound by the procedure of example 32. LC MS m/z = 437, (MH<sup>+</sup>), Rt = 1.44 min.

Example 74

1-(2,6-difluorophenyl)-5-phenyl-7-(4-carboxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

15 The product of example 60 was converted to the target compound by the procedure of example 34. LC MS m/z = 466 (MH<sup>+</sup>), Rt = 2.39 min

Example 75

1-(2,6-difluorophenyl)-5-(4-fluorophenyl)-7-(4-carboxypiperidin-1-yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

20 The product of example 67 was converted to the target compound by the procedure of example 34. LC MS m/z = 484 (MH<sup>+</sup>), Rt = 1.95 min

Example 76

1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

25 The product of example 50 (75 mg, 0.168 mmol), pyrrolidin-1-yl ethylamine (57mg, 0.5 mmol) were combined in a sealed tube at 90° for 20 h. Concentration and prep hplc afforded the title compound as an amber oil. LC MS m/z = 480 (MH<sup>+</sup>), Rt = 1.52 min

Example 77

30 1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1H-tetrazol-5-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one

The method of example 76 was repeated except using 5-aminotetrazole as the amine and the reaction was heated to 150° for 10 h. Concentration and prep hplc afforded the title compound as an amber oil. LC MS m/z = 450 (MH<sup>+</sup>), Rt = 1.94 min.

35 The compounds in **Table 6** were prepared by either the method of Example 76 (method G) or the method of Example 77 (method H) using the appropriate amine and the sulphone of the example indicated.

Table 6

Example	Compound name	Method*	LC MS	
			m/z	Rt min
78	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-ylethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	495	1.63
79	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1-hydroxymethyl-1-methylethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) H	472	1.45
80	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methylethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	442	1.62
81	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methylethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	442	1.60
82	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) H	488	1.31
83	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino) -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) H	447	2.26
84	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(2-hydroxyethyl)-methylamino] -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	442	1.75
85	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(racemic)-3-exo-(bicyclo[2.2.1]hept-ene-2-carboxylic acid amide)amino] -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) H	519	1.87
86	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(racemic)-3-endo-(bicyclo[2.2.1]hept-ene-2-carboxylic acid amide)amino] -3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	519	1.80

Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
87	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7- {([1,3]dioxolan-2-ylmethyl)-amino}-3,4-dihydro- 1H-pyrimido[4,5-d]pyrimidin-one	(50) G	470	1.77
88	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7- [3-(2-oxo-pyrrolidin-1-yl) propylamino]-3,4- dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) G	509	1.69
89	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7- (2-hydroxyethylamino) -3,4-dihydro-1H- pyrimido[4,5-d]pyrimidin-one	(50) G	427	1.53
90	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-(2-pyrrolidin-1-yl-ethylamino) -3,4-dihydro-1H- pyrimido[4,5-d]pyrimidin-one	(49) G	475	1.59
91	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-(2-piperidin-1-yl ethylamino) -3,4-dihydro-1H- pyrimido[4,5-d]pyrimidin-one	(49) G	489	1.65
92	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-(1H-tetrazol-5-ylamino) -3,4-dihydro-1H- pyrimido[4,5-d]pyrimidin-one	(49) H	446	2.10
93	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-((R)-2-hydroxy-1-methyl-ethylamino) -3,4- dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) G	436	1.69
94	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-[(2-hydroxyethyl)-methylamino] -3,4-dihydro- 1H-pyrimido[4,5-d]pyrimidin-one	(49) G	436	1.79
95	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-[(1,3]dioxolan-2-ylmethyl)amino]-3,4-dihydro- 1H-pyrimido[4,5-d]pyrimidin-one	(49) G	464	1.85
96	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)- 7-[3-(2-oxo-pyrrolidin-1-yl)-propylamino]-3,4- dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) G	503	1.80

Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
97	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) G	422	1.63
98	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)-7-(3-pyrrolidin-1-ylpropylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) G	489	1.47
99	5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-ylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	453	1.55
100	5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-ylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	467	1.72
101	5-(4-fluoro-2-methylphenyl)-7-(1H-tetrazol-5-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) H	424	2.17
102	5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1-hydroxymethyl-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	444	1.66
103	5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	414	1.68
104	5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	428	2.06
105	5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) H	460	1.52
106	5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	428	1.85



Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
107	5-(4-fluoro-2-methylphenyl)-7-[(1,3)dioxolan-2-ylmethyl)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	442	1.84
108	5-(4-fluoro-2-methylphenyl)-7-[3-(2-oxo-pyrrolidin-1-yl)-propylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	481	1.72
109	5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	400	1.59
110	5-(4-fluoro-2-methylphenyl)-7-(3-pyrrolidin-1-ylpropylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) G	467	1.47
111	1-(2-Methylphenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	461	1.42
112	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	475	1.49
113	1-(2-Methylphenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1-hydroxymethyl-1-methyl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	452	1.41
114	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	421	1.53
115	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	422	1.54
116	1-(2-Methylphenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) H	468	1.30

Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
117	-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	436	1.60
118	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[(2-hydroxyethyl)-methylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	422	1.66
119	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[[1,3]dioxolan-2-ylmethyl)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	450	1.70
120	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[3-(2-oxo-pyrrolidin-1-yl)-propylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	489	1.60
121	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) G	408	1.42
122	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	475	1.48
123	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	489	1.55
124	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(1H-tetrazol-5-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) H	446	1.95
125	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1-hydroxymethyl-1-methyl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) H	466	1.49

Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
126	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	436	1.58
127	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	436	1.60
128	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) H	482	1.36
129	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	450	1.70
130	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[(2-hydroxyethyl)-methylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	436	1.72
131	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[[1,3]dioxolan-2-ylmethyl)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	464	1.77
132	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-[3-(2-oxo-pyrrolidin-1-yl)-propylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	503	1.68
133	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-piperazin-1-yl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G <sup>a</sup>	447	2.76

Table 6 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
134	1-(2,6-Dimethylphenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	422	1.51
135	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(3-pyrrolidin-1-ylpropylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) G	489	1.44
136	1-(2-Fluorophenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	465	1.49
137	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-yl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	479	1.53
138	1-(2-Fluorophenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1-hydroxymethyl-1-methyl ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) H	456	1.45
139	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	426	1.54
140	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methylethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	440	1.78
141	1-(2-Fluorophenyl)- 5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxymethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) H	472	1.34
142	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	440	1.65

Table 65 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
143	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(2-hydroxyethyl)-methylamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	426	1.70
144	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(racemic)-3-exo-(bicyclo[2.2.1]hept-ene-2-carboxylic acid amide)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) H	503	1.82
145	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[(racemic)-3-endo-(bicyclo[2.2.1]hept-ene-2-carboxylic acid amide)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) H	503	1.79
146	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[[1,3]dioxolan-2-ylmethyl)amino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	454	1.76
147	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) G	412	1.46

\* the sulfone product of the exemplified example number;

a. The reactant amine was 1-Boc-piperazine. The Boc group in this case was removed during the reaction.

5

#### Example 148

1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-ethylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one

The product of example 52 (75 mg, 0.17 mmol), 1-ethylpiperidin-4-ylamine hydrochloride (100 mg, 0.5 mmol), K<sub>2</sub>CO<sub>3</sub> (83 mg, 0.6 mmol) and NMP (1 mL) were combined and heated to 150° in a sealed tube for 10 h. Concentration and prep hplc afforded the title compound as an amber oil. LC MS m/z = 475 (MH<sup>+</sup>), Rt = 1.42 min

#### Example 149

1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one

The product of example 54 (75 mg, 0.17 mmol), 1-methylpiperidin-4-ylamine hydrochloride (94 mg, 0.5 mmol), K<sub>2</sub>CO<sub>3</sub> (83 mg, 0.6 mmol) and NMP (1 mL) were combined and heated to 90° in a sealed tube for 10h. Concentration and prep hplc

afforded the title compound as an amber oil. LC MS  $m/z$  = 465 ( $MH^+$ ),  $R_t$  = 1.49 min.

The products of the examples in **Table 7** are made by either the method of example 148 (Method I) or the method of example 149 (Method J) using the sulfone product from **Table 4** as indicated and the appropriate amine hydrochloride.

**Table 7**

Example	Compound name	Method*	LC MS	
			$m/z$	$R_t$ min
150	1-(2-Chlorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) J	481	1.44
151	1-(2-Chlorophenyl)-5-(4-fluoro-2-ethylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(50) I	495	1.50
152	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) J	475	1.48
153	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)-7-(1-carboxyethyl-2-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) J	436	2.20
154	1-((S)-1-Phenylethyl)-5-(4-fluoro-2-methylphenyl)-7-(1-carboxyethyl-2-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(49) J	449	1.64
155	1-Cyclohexyl-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) $J^a$	453	1.47
156	1-Cyclohexyl-5-(4-fluoro-2-methylphenyl)-7-(1-ethylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(51) I	467	1.54
157	1-(2-Methylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(52) J	461	1.37

Table 7 cont.

Example	Compound name	Method*	LC MS	
			m/z	Rt min
158	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) J	475	1.42
159	1-(2,6-Dimethylphenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-ethylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(53) J	489	1.47
160	1-(2-Fluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-ethylpiperidin-4-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-one	(54) I	479	1.49

\* the sulfone product of the exemplified example number

a. The starting amine is glycine tert-butyl ester. The ester is cleaved under the reaction conditions.

5

#### Example 161

##### 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of Example 3 (0.102 g, 0.22 mmol) was suspended in MeOH (2 mL) and stirred under argon. A 1 Molar solution of NaOMe in methanol (0.44 mL, 0.44 mmol) was added. After 10 min the solvent was removed *in vacuo*, and the residue partitioned between EtOAc and H<sub>2</sub>O. The organic phase was washed with H<sub>2</sub>O (2 X), satd aq NaCl (1 X), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give the crude product. Recrystallization from EtOAc/hexane gave the title compound as a white-crystalline solid. mp 210-211°C, LC MS m/z = 401 (MH<sup>+</sup>) Rt = 2.0 min.

15

#### Example 162

##### 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-ethoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of Example 3 (0.0658 g, 0.147 mmol) was suspended in EtOH (2 mL) and stirred under argon. A 0.5 M solution of NaOEt in EtOH (0.587 mL, 0.294 mmol) was added. After 15 min the solvent was removed *in vacuo*, and the residue partitioned between EtOAc and H<sub>2</sub>O. The organic phase was washed with H<sub>2</sub>O (2 X), satd aq NaCl (1 X), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give the crude product. Flash chromatography on silica gel eluted with 0-5% EtOAc/CH<sub>2</sub>Cl<sub>2</sub> gave the title compound as a white-amorphous solid. mp 191-194°C, LC MS m/z = 415

25

(MH+) Rt = 2.15 min.

#### Example 163

5 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethoxy)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

The product of Example 3 (0.0658 g, 0.147 mmol) was suspended in ethylene glycol (1 mL) and stirred under argon. NaH, dry, 95%, (8.0 mg, 0.32 mmol) was added. After 30 min the solvent was removed *in vacuo*, and the residue partitioned between EtOAc and H<sub>2</sub>O. The organic phase was washed with H<sub>2</sub>O (2 X), satd aq NaCl (1 X),  
10 dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give the crude product. Flash chromatography on silica gel eluted with 0-10% EtOAc/CH<sub>2</sub>Cl<sub>2</sub> gave the title compound as a white-amorphous solid. mp 172-175°C, LC MS m/z = 431 (MH+) Rt = 1.7 min.

#### Example 164

15 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-yloxy)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one trifluoroacetate

A solution of 4-hydroxy-1-methylpiperidine (35.1 mg, 0.3 mmol) in DMSO (0.25 ml) was stirred under argon and NaH, dry, 95%, (7.5 mg, 0.3 mmol) was added. The mixture was stirred for 15 min, and then a solution of the product of Example 3 (0.0672 g, 0.15 mmol) dissolved in DMSO (0.25 mL) was added. After the mixture was stirred  
20 for 30 min, the reaction mixture was partitioned between EtOAc and H<sub>2</sub>O. The organic phase was washed with H<sub>2</sub>O (5 X), satd aq NaCl (1 X), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give the crude product. Purification by preparative hplc followed by lyophilization gave the title compound as a white-amorphous solid. LC MS  
25 m/z = 484 (MH+) Rt = 1.47 min.

#### Example 165

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[[bis-(2-hydroxyethyl)amino]ethoxy]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one  
30 trifluoroacetate

NaH, dry, 95%, (8 mg, 0.4 mmol) was added to triethanolamine (243 mg, 1.6 mmol), and then DMSO (0.25 ml) was added. The mixture agitated for 2 min., and then the product of Example 3 (0.0896 g, 0.2 mmol) was added. After 5 min of agitation, additional DMSO (0.25 mL) was added. After the mixture was agitated for 10 min, the  
35 reaction mixture was partitioned between EtOAc and H<sub>2</sub>O. The organic phase was washed with H<sub>2</sub>O (5 X), satd aq NaCl (1 X), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give the crude product. Purification by preparative hplc followed by



lyophilization gave the title compound as a white-amorphous solid. LC MS  $m/z$  = 518 (MH<sup>+</sup>)  $R_t$  = 1.25 min.

Example 166

5    1-(2,6-Difluorophenyl)-5-phenyl-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

Following the procedure of example 161 except using the product of example 45 in place of the product of example 3 gave the title compound as a white-crystalline solid. mp 221-224°C, LC MS  $m/z$  = 369 (MH<sup>+</sup>)  $R_t$  = 1.92 min.

10

Example 167

1,5-diphenyl-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

a)    1-(2,6-difluorophenyl)-5-phenyl-7-methylsulfonyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

15    Following the procedure of example 3 except for using the product of example 1 in place of the product of example 2 gave the title compound as a white solid. LC MS  $m/z$  = 381 (MH<sup>+</sup>)  $R_t$  = 1.78 min.

b)    1,5-diphenyl-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one

20    Following the procedure of example 161 except using the product of example 167 (a) in place of the product of example 3 gave the title compound as a white-amorphous solid. mp 220-222°C, LC MS  $m/z$  = 333 (MH<sup>+</sup>)  $R_t$  = 1.94 min.


25    All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

30    The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore, the Examples herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is Claimed is:

1. A compound of the formula:

5



wherein

R<sub>1</sub> is a aryl or a heteroaryl ring, which ring is optionally substituted;

R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, aryl,  
 10 arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a  
 heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties,  
 excluding hydrogen, are optionally substituted;

R<sub>3</sub> is a C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>alkyl, aryl,  
 arylC<sub>1-10</sub>alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or a  
 15 heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are  
 optionally substituted;

Y is a bond, CR<sub>b</sub>, C(O), N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>;

R<sub>b</sub> is hydrogen, C<sub>1-2</sub> alkyl, NR<sub>c</sub>, hydroxy, thio, C<sub>1-2</sub> alkoxy, or S(O)<sub>m</sub>C<sub>1-2</sub> alkyl;

R<sub>c</sub> is hydrogen or C<sub>1-2</sub> alkyl;

20 R<sub>d</sub> is hydrogen or C<sub>1-2</sub> alkyl;

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

m is 0 or an integer having a value of 1 or 2;

n is 0 or an integer having a value of 1 to 10;

R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted  
 25 C<sub>1-6</sub> alkyl, optionally substituted aryl or an optionally substituted aryl-C<sub>1-4</sub>  
 alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a  
 heterocyclic ring of 5 to 7 members which ring optionally contains an additional  
 heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and wherein the ring is  
 optionally substituted;  
 30 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl,  
 aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety; and wherein  
 each of these moieties, excluding hydrogen, is optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted arylC<sub>1-4</sub> alkyl;

Z is oxygen or sulfur;

or a pharmaceutically acceptable salt thereof.

5

2. The compound according to Claim 1 wherein the structure is of Formula (IV).

3. The compound according to Claim 1 wherein R<sub>3</sub> is an optionally substituted aryl, arylC<sub>1-10</sub>alkyl, or a cycloalkyl moiety.

10

4. The compound according to Claim wherein R<sub>1</sub> is an optionally substituted aryl moiety.

15

5. The compound according to any one of Claims 3 or 4 wherein the moiety is optional substituted one or more times, independently by halogen, alkyl, hydroxy, alkoxy, amino, or halosubstituted alkyl.

6. The compound according to any one of Claims 1 to 5 wherein X is OR<sub>2</sub>.

20

7. The compound according to any one of Claims 1 to 5 wherein X is S(O)<sub>m</sub>R<sub>2</sub>.

8. The compound according to any one of Claims 1 to 5 wherein X is (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>.

25

9. The compound according to any one of Claims 1 to 5 wherein X is (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>.

10. The compound according to any one of Claims 1 to 5 wherein X is R<sub>2</sub>.

30

11. The compound according to any one of Claims 1 to 10 wherein Y is CR<sub>b</sub>, or C(O).

12. The compound according to any one of Claims 1 to 10 wherein Y is N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>.

35

13. The compound according to any one of Claims 1 to 10 wherein Y is a bond.

14. The compound according to any one of Claims 1 to 13 wherein R<sub>2</sub> is an

optionally substituted alkyl, heteroaryl, or heterocyclic or heterocyclic alkyl.

15. The compound according to Claim 14 wherein R<sub>2</sub> is an optionally substituted imidazole, tetrazole, pyrrolidine, piperidine, piperazine, or morpholine ring.

5

16. The compound according to Claim 15 wherein the ring is substituted one or more times, independently, by halogen, C<sub>1-10</sub>alkyl, halosubstituted C<sub>1-10</sub>alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>.

10

17. The compound according to Claim 1 which is

7-Methylsulfanyl-1,5-diphenyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfanyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-methylsulfonyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

15

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-4-ylamino)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one trifluoroacetate;

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-1,3-dihydroxyprop-2-ylamino)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

20

5-(4-Fluoro-2-methylphenyl)-7-methylsulfanyl-1-((*R*)-1-phenylethyl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

5-(4-Fluoro-2-methylphenyl)-7-methylsulfanyl-1-((*S*)-1-phenylethyl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperazin-4-yl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

25

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperazin-4-yl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methyl-1*H*-imidazol-2-yl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

30

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(imidazol-1-yl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

1-(2,6-Difluorophenyl)-7-methylsulfanyl-5-phenyl-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

1-(2,6-Difluorophenyl)-7-methylsulfanyl-5-(4-fluorophenyl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]pyrimidin-2-one;

35

1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-([2-hydroxyethyl])-

- methlamino]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-piperazin-4-yl)-3,4-  
dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-hydroxypiperidin-1-yl)-  
5 3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-carboxypiperidin-1-yl)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-carboethoxypiperidin-1-  
yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
10 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(4-hydroxymethylpiperidin-1-  
yl)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-di(ethan-2-ol-)amino-3,4-  
dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-yl-  
15 amino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-morpholin-4-yl)-3,4-  
dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-morpholin-4-yl ethylamino)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
20 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyrrolidin-1-yl ethylamino)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-piperidin-1-yl ethylamino)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one; SB-647851-M  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyridin-3-yl ethylamino)-  
25 3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-diethylamino ethylamino)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-pyridin-3-yl ethylamino)-  
3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;  
30 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2- dimethylamino)-3,4-  
dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one ;  
1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2- methylamino)-3,4-dihydro-  
1H-pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(2-methylphenyl)-7-methylsulfanyl-3,4-dihydro-1H-  
35 pyrimido[4,5-d]pyrimidin-2-one;  
1-(2,6-Difluorophenyl)-5-(2-methylphenyl)-7-(1-piperazin-4-yl)-3,4-dihydro-1H-  
pyrimido[4,5-d]pyrimidin-2-one;

- 1-(2,6-Difluorophenyl)-5-(2-methylphenyl)-7-(2-diethylamino ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-phenyl-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 5 1-(2-Fluorophenyl)-7-methylsulfanyl-5-(2-methyl-4-fluoro)phenyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Dimethylphenyl)-7-methylsulfanyl-5-(2-methyl-4-fluoro)phenyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2-Methylphenyl)-7-methylsulfanyl-5-(2-methyl-4-fluoro)phenyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 10 1-Cyclohexyl-7-methylsulfanyl-5-(2-methyl-4-fluoro)phenyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2-Chlorophenyl)-7-methylsulfanyl-5-(2-methyl-4-fluoro)phenyl-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 15 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-methyl-1,3-dihydroxyprop-2-ylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-ethoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((R)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 20 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-((S)-2-hydroxy-1-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-dimethyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 25 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxy-1,1-bis-hydroxy-methyl-ethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(1-methylpiperidin-4-yloxy)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 30 1,5-Diphenyl-7-methoxy-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2-hydroxyethoxy)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-[[bis-(2-hydroxyethyl)-amino]ethoxy]-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one;
- 35 1-(2,6-Difluorophenyl)-5-(4-fluoro-2-methylphenyl)-7-(2,3-dihydroxypropylamino)-3,4-dihydro-1H-pyrimido[4,5-d]pyrimidin-2-one; or

a pharmaceutically acceptable salt thereof.

18. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 17, and a pharmaceutically acceptable carrier or diluent.


5

19. A method of treating, including prophylaxis, of a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a compound of according to any one of Claims 1 to 17.

10 20. The method according to Claim 19 wherein the CSBP/RK/p38 kinase mediated disease is psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, meningitis, ischemic and hemorrhagic stroke,  
15 neurotrauma/closed head injury, asthma, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, chronic obstructive pulmonary disease, silicosis, pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, cardiac and brain and renal reperfusion injury, congestive heart failure, coronary arterial bypass grafting (CABG) surgery, thrombosis, glomerularnephritis, chronic renal failure, diabetes, diabetic retinopathy, macular degeneration, graft vs. host reaction, allograft rejection,  
20 inflammatory bowel disease, Crohn's disease, ulcerative colitis, neurodegenerative disease, muscle degeneration, diabetic retinopathy, macular degeneration, tumor growth and metastasis, angiogenic disease, influenza induced pneumonia, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis.

25

21. A compound of the formula :



30 wherein

R1 is an aryl or heteroaryl ring, which ring is optionally substituted;

R2 is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylalkyl, aryl, arylC<sub>1-10</sub>

- alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, are optionally substituted;
- R<sub>3</sub> is C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkyl C<sub>1-10</sub> alkyl, arylC<sub>1-10</sub> alkyl, heteroarylC<sub>1-10</sub> alkyl, heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are optionally substituted;
- X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;
- n is 0 or an integer having a value of 1 to 10;
- m is 0 or an integer having a value of 1 or 2;
- R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl or an optionally substituted aryl-C<sub>1-4</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>;
- R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl; and wherein each of these moieties, excluding hydrogen, may be optionally substituted;
- R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;
- Z is oxygen or sulfur;  
or a pharmaceutically acceptable salt thereof.
22. The compound according to Claim 21 wherein the structure is of Formula (I).
23. The compound according to Claim 22 wherein R<sub>1</sub> and/or R<sub>3</sub> are an optionally substituted aryl.
24. The compound according to Claim 23 wherein the aryl is substituted one or more times, independently, by halogen, alkyl, hydroxy, alkoxy, amino, or halosubstituted alkyl.
25. The compound according to any one of Claims 21 to 24 wherein X is R<sub>2</sub>, OR<sub>2</sub>, or S(O)<sub>m</sub>R<sub>2</sub>.
26. The compound according to any one of Claims 21 to 24 wherein X is (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>.



27. The compound according to Claim 21, which is:


7-Methylsulfonyl-5-phenyl-1-(1-phenylethyl)-3,4-dihydro-1*H*-pyrimido[4,5-*d*]-pyrimidin-2-one; or a pharmaceutically acceptable salt thereof.

5 28. A pharmaceutical composition comprising a compound according to any one of Claims 21 to 27, and a pharmaceutically acceptable carrier or diluent.

29. A method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount  
10 of a compound of Formula (I) or (Ia) according to any one of Claims 21 to 27.

30. The method according to Claim 29 wherein the CSBP/RK/p38 kinase mediated disease is psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis  
15 and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, meningitis, ischemic and hemorrhagic stroke, neurotrauma/closed head injury, asthma, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, chronic obstructive pulmonary disease, silicosis, pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, cardiac and brain  
20 and renal reperfusion injury, congestive heart failure, coronary arterial bypass grafting (CABG) surgery, thrombosis, glomerulonephritis, chronic renal failure, diabetes, diabetic retinopathy, macular degeneration, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, neurodegenerative disease, muscle degeneration, diabetic retinopathy, macular degeneration, tumor growth and  
25 metastasis, angiogenic disease, influenza induced pneumonia, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis.

31. A compound of the formula:



30

wherein

- R<sub>1</sub> is an aryl or heteroaryl ring, which ring is optionally substituted;  
R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylalkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, are  
5 optionally substituted;  
R<sub>3</sub> is an C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylalkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties are optionally substituted;  
Y is CR<sub>b</sub>, C(O), N(R<sub>d</sub>), oxygen, or S(O)<sub>m</sub>;  
10 R<sub>b</sub> is hydrogen, C<sub>1-2</sub> alkyl, NR<sub>c</sub>, hydroxy, thio, C<sub>1-2</sub> alkoxy, S(O)<sub>m</sub>C<sub>1-2</sub> alkyl,  
R<sub>c</sub> is hydrogen or C<sub>1-2</sub> alkyl;  
R<sub>d</sub> is hydrogen or C<sub>1-2</sub> alkyl;  
X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;  
m is 0 or an integer having a value of 1 or 2;  
15 n is 0 or an integer having a value of 1 to 10;  
R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl or an optionally substituted aryl-C<sub>1-4</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional  
20 heteroatom selected from oxygen, sulfur or NR<sub>9</sub>;  
R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl; and wherein each of these moieties, excluding hydrogen, may be optionally substituted;  
R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub> or optionally substituted C<sub>1-10</sub> alkyl, optionally substituted  
25 aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;  
Z is oxygen or sulfur;  
or a pharmaceutically acceptable salt thereof.

32. The compound according to Claim 31 wherein the structure is of Formula (II).

30

33. The compound according to Claim 31 wherein R<sub>1</sub> and/or R<sub>3</sub> is an optionally substituted aryl.

34. The compound according to Claim 33 wherein the aryl is substituted one or more  
35 time, independently, by halogen, alkyl, hydroxy, alkoxy, amino, or halosubstituted alkyl.

35. The compound according to Claim 31 wherein X is R<sub>2</sub>, OR<sub>2</sub>, or S(O)<sub>m</sub>R<sub>2</sub>.

36. The compound according to Claim 31 wherein X is or  $(CH_2)_n NR_4R_{14}$ , or  $(CH_2)_n NR_2R_4$ .

37. The compound according to any one of Claims 31 to 36 wherein  $R_2$  is an optionally substituted heteroaryl or heterocyclic.

38. A pharmaceutical composition comprising a compound according to any one of Claims 31 to 37, and a pharmaceutically acceptable carrier or diluent.


10

39. A method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a compound of Formula (II) or (IIa) according to any one of Claims 31 to 37.


40. The method according to Claim 39 wherein the CSBP/RK/p38 kinase mediated disease is psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, meningitis, ischemic and hemorrhagic stroke, neurotrauma/closed head injury, asthma, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, chronic obstructive pulmonary disease, silicosis, pulmonary sarcosis, bone resorption disease, osteoporosis, restenosis, cardiac and brain and renal reperfusion injury, congestive heart failure, coronary arterial bypass grafting (CABG) surgery, thrombosis, glomerulonephritis, chronic renal failure, diabetes, diabetic retinopathy, macular degeneration, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, neurodegenerative disease, muscle degeneration, diabetic retinopathy, macular degeneration, tumor growth and metastasis, angiogenic disease, influenza induced pneumonia, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis.

30

41. A compound of the formula:



(III) or



(IIIa)

wherein

R<sub>1</sub> is an aryl, or a heteroaryl ring, which ring is optionally substituted;

R<sub>2</sub> is a heterocyclic or heterocyclic C<sub>1-10</sub> alkyl ring, which is ring is optionally substituted;

5 R<sub>2</sub>' is a heterocyclic, heterocyclic C<sub>1-10</sub> alkyl, aryl, or heteroaryl ring, which ring is optionally substituted;

R<sub>3</sub> is an aryl or heteroaryl ring, which ring is optionally substituted;

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub> NR<sub>2</sub>R<sub>4</sub>; provided that when X is (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, then R<sub>4</sub>R<sub>14</sub> must form the optionally

10 substituted cyclized ring optionally comprising an oxygen, sulfur or nitrogen moiety;

n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

R<sub>4</sub> and R<sub>14</sub> are independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl, or optionally substituted arylC<sub>1-4</sub> alkyl; or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>, and which ring is optionally substituted;

15 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or a heteroarylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen, is optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl, or optionally substituted arylC<sub>1-4</sub> alkyl;

Z is oxygen or sulfur;

25 or a pharmaceutically acceptable salt thereof.


42. The compound according to Claim 41 wherein the structure is of Formula (III).

43. The compound according to Claim 41 wherein R<sub>1</sub> and/or R<sub>3</sub> is an optionally substituted aryl.

44 The compound according to Claim 43 wherein the aryl is substituted one or more times, independently by halogen, alkyl, hydroxy, alkoxy, amino, or halosubstituted alkyl.


35 45. The compound according to Claim 42 wherein X is OR<sub>2</sub>, or S(O)<sub>m</sub>R<sub>2</sub>.

46. The compound according to Claim 42 wherein X is (CH<sub>2</sub>)<sub>n</sub> NR<sub>4</sub>R<sub>14</sub>, or




47. The compound according to Claim 42 wherein X is R<sub>2</sub>'.
- 5 48. The compound according to Claim 47 wherein R<sub>2</sub>' is an optionally substituted heteroaryl or heterocyclic ring.
49. The compound according to Claim 48 wherein the ring is an optionally substituted imidazole, pyrrolidine, piperidine, piperazine, or a morpholine ring.
- 10 50. The compound according to Claim 49 wherein the ring is substituted one or more times, independently, by halogen, C<sub>1-10</sub>alkyl, halosubstituted C<sub>1-10</sub>alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>6</sub>, and (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>C(Z)OR<sub>6</sub>.
- 15 51. A pharmaceutical composition comprising a compound according to any one of Claims 41 to 50, and a pharmaceutically acceptable carrier or diluent.
52. A method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount
- 20 of a compound of Formula (III) or (IIIa) according to any one of Claims 41 to 50.
53. The method according to Claim 52 wherein the CSBP/RK/p38 kinase mediated disease is psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis,
- 25 and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, meningitis, ischemic and hemorrhagic stroke, neurotrauma/closed head injury, asthma, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, chronic obstructive pulmonary disease, silicosis, pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, cardiac and brain
- 30 and renal reperfusion injury, congestive heart failure, coronary arterial bypass grafting (CABG) surgery, thrombosis, glomerular nephritis, chronic renal failure, diabetes, diabetic retinopathy, macular degeneration, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, neurodegenerative disease, muscle degeneration, diabetic retinopathy, macular degeneration, tumor growth and
- 35 metastasis, angiogenic disease, influenza induced pneumonia, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis.

54. A compound of the formula:



(V) or



(Va)

5 wherein

X is R<sub>2</sub>, OR<sub>2</sub>, S(O)<sub>m</sub>R<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>NR<sub>4</sub>R<sub>14</sub>, or (CH<sub>2</sub>)<sub>n</sub>NR<sub>2</sub>R<sub>4</sub>;

n is 0 or an integer having a value of 1 to 10;

m is 0 or an integer having a value of 1 or 2;

R<sub>1</sub> is an optionally substituted aryl ring;

10 R<sub>2</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkylalkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, heterocyclic, or heterocyclylC<sub>1-10</sub> alkyl moiety; and wherein each of these moieties, excluding hydrogen are optionally substituted;

R<sub>3</sub> is an optionally substituted aryl ring;

15 R<sub>4</sub> and R<sub>14</sub> are each independently selected from hydrogen, optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl, or R<sub>4</sub> and R<sub>14</sub> together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub>;

20 R<sub>6</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl; and wherein each of these moieties, excluding hydrogen, may be optionally substituted;

R<sub>9</sub> is hydrogen, C(Z)R<sub>6</sub>, optionally substituted C<sub>1-10</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;

25 Z is oxygen or sulfur;


or a pharmaceutically acceptable salt thereof.

55. The compound according to Claim 54 wherein R<sub>1</sub> and R<sub>3</sub> are optionally substituted phenyl rings.


30

56. The compound according to Claim 55 wherein the aryl is substituted one or more times, independently, by halogen, alkyl, hydroxy, alkoxy, amino, or halosubstituted alkyl.


57. The compound according to Claim 56 wherein  $R_3$  is unsubstituted or substituted one or more times, independently with halogen.
- 5 58. The compound according to Claim 55, wherein  $R_1$  is unsubstituted or substituted one or more times, independently with halogen or alkyl.
59. The compound according to any one of Claims 54 to 58 wherein X is  $OR_2$ .
- 10 60. The compound according to any one of Claims 54 to 58 wherein X is  $S(O)_mR_2$ .
61. The compound according to any one of Claims 54 to 58 wherein X is  $(CH_2)_nNR_4R_{14}$ .
- 15 62. The compound according to any one of Claims 54 to 58 wherein X is  $(CH_2)_nNR_2$ .
63. The compound according to any one of Claims 54 to 58 wherein X is  $R_2$ .
- 20 64. The compound according to Claim 54 wherein  $R_2$  is an optionally substituted alkyl, heteroaryl, or heterocyclic moiety.
65. The compound according to Claim 64 wherein  $R_2$  is an optionally substituted imidazole, pyrrolidine, piperidine, piperazine, or a morpholine ring.
- 25 66. The compound according to Claim 65 wherein the ring is substituted one or more times, independently, by halogen,  $C_{1-10}$ alkyl, halosubstituted  $C_{1-10}$ alkyl,  $(CR_{10}R_{20})_nNR_4R_{14}$ ,  $(CR_{10}R_{20})_nOR_6$ , and  $(CR_{10}R_{20})_nC(Z)OR_6$ .
- 30 67. The compound according to any one of Claims 54 to 66 wherein the structure is of Formula (V).
68. A process for producing a compound of Formula (IV) according to Claim 1, which process comprises reacting a compound of the formula
- 35



wherein Y-R<sub>1</sub> and R<sub>3</sub> are as defined herein for Formula (IV); and X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl or heteroaryl moiety; under cyclization conditions to yield a compound of the formula:




wherein Y-R<sub>1</sub> and R<sub>3</sub> are as defined herein for Formula (IV), and X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl or heteroaryl moiety; and if necessary displacing the sulfoxide or sulfone of Formula (B) with a compound of the formula:



wherein R<sub>2</sub> is as is as defined for Formula (IV); and X is O/N/S; to yield a compound of Formula (IV).

69. The process according to Claim 68 wherein the compound of Formula (A) is produced by reacting a compound of the formula:



wherein R<sub>1</sub> and R<sub>3</sub> are as defined in Formula (IV), and X<sub>1</sub> is a C<sub>1-10</sub> alkyl, aryl or heteroaryl moiety; with a suitable reducing agent to yield a compound of Formula (A).



## p38/CSBP Kinase Pathway




FIGURE 1

FIGURE 2

## Citric Acid Induced Cough Model




FIGURE 3

# Antigen- or LTD<sub>4</sub>-Induced Hypertussive Model in the Guinea Pig





Figure 4

## Effects of Dextromethorphan or Codeine On Citric Acid-Induced Cough in Guinea Pigs



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/06688

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 487/04; A61K 31/519

US CL : 544/256; 514/258

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbol)

U.S. : 544/256; 514/258

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4,886,807 A (KITAMURA et al) 12 December 1989 (12.12.1989), see formula (I).	1-5, 17, 21-27, 31-37, 41-66
A, P	US 6,150,373 A (HARRIS et al) 21 November 2000 (21.11.2000), see formula (I).	1-5, 17, 21-27, 31-37, 41-66 and 68

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

05 April 2001 (05.04.2001)

Date of mailing of the international search report

06 JUN 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Deepak Rao

Telephone No. (703) 308-1235

JOYCE BRIDGERS  
PARALEGAL SPECIALIST  
CHEMICAL MATRIN

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/06688

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 6-16, 18-20, 28-30, 38-40 and 67  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐  
☐

- The additional search fees were accompanied by the applicant's protest.  
No protest accompanied the payment of additional search fees.